

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number  
WO 01/83459 A2

(51) International Patent Classification<sup>7</sup>: C07D 239/00

(21) International Application Number: PCT/EP01/04850

(22) International Filing Date: 30 April 2001 (30.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/201,824 4 May 2000 (04.05.2000) US

(71) Applicant: BASF AKTIENGESELLSCHAFT  
[DE/DE]; 67056 Ludwigshafen (DE).

(72) Inventors: CARLSEN, Marianne; 222 Arborlea Avenue, Yardley, PA 19067 (US). GUACIARO, Michael, Anthony; 21 Inverness Lane, East Windsor, NJ 08520 (US). TAKASUGI, James, Jan; 11 Pembroke Court, Lawrenceville, NJ 08648 (US).

(74) Common Representative: BASF AKTIENGESELLSCHAFT; 67056 Ludwigshafen (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: URACIL SUBSTITUTED PHENYL SULFAMOYL CARBOXAMIDES



(57) Abstract: Novel uracil substituted phenyl sulfamoyl carboxamides (I) and salts thereof, where A = oxygen or sulfur; X<sup>1</sup> = H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl; X<sup>2</sup> = H, CN, CS-NH<sub>2</sub>, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl; X<sup>3</sup> = H, CN, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxyalkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, optionally substituted benzyl; R<sup>1</sup>, R<sup>2</sup> = H, halogen, optionally substituted hydroxy, C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, C<sub>3</sub>-C<sub>10</sub>-alkynyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, phenyl, benzyl or C<sub>5</sub>-C<sub>7</sub>-cycloalkenyl, or R<sup>1</sup> + R<sup>2</sup> together with the atom to which they are attached form a 3- to 7-membered heterocyclic

ring; Q is selected from Q<sup>1</sup> to Q<sup>40</sup> as defined in the description. Use: as herbicides; for the desiccation/defoliation of plants.

WO 01/83459 A2

## URACIL SUBSTITUTED PHENYL SULFAMOYL CARBOXAMIDES

## Description

5

Weeds cause tremendous global economic losses by reducing crop yields and lowering crop quality. Worldwide, agronomic crops must compete with hundreds of weed species.

10 In spite of the commercial herbicides available today, damage to crops caused by weeds still occurs. Accordingly, there is ongoing research to create more effective and/or more selective herbicidal agents.

In WO 98/06706 are disclosed the use of certain p-trifluoro-  
15 methylphenyl uracils, their method of production and their use as herbicides. In addition, WO 96/08151 discloses herbicidal aryl uracils and arylthiouracils in which the aryl ring is an optionally substituted phenyl group. In neither disclosure, however is there mentioned a sulfamoyl carboxamide group  
20 substituent.

Therefore, it was an object of the present invention to provide novel 3-phenyluracils which are highly effective for the control of undesirable plant species. The object also extends to  
25 providing novel compounds which act as desiccants/defoliants.

It was also an object of the present invention to provide a method for the control of undesirable plant species and compositions useful therefor.


30

It is an advantage of the present invention that the method for the control of undesirable plant species may be employed in the presence of a crop.

35 It was a further object of the present invention to provide a process for the preparation of herbicidal phenylsulfamoyl carboxamides and an intermediate compound useful therefor.

These and other objects and advantages of the present invention  
40 will become more apparent from the detailed description thereof set forth below.

We have found that this object is achieved in accordance with the invention by the novel uracil substituted phenylsulfamoyl carb-  
45 oxamides of the formula I



I,

wherein the variables have the following meanings:

A oxygen or sulfur;

$X^1$  hydrogen, halogen or  $C_1$ - $C_4$ -alkyl;

$X^2$  hydrogen, cyano,  $CS-NH_2$ , halogen,  $C_1$ - $C_4$ -alkyl or  $C_1$ - $C_4$ -haloalkyl;





$X^3$  hydrogen, cyano,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkoxy-alkyl,  $C_3$ - $C_7$ -cycloalkyl,  $C_3$ - $C_6$ -alkenyl,  $C_3$ - $C_6$ -alkynyl or optionally substituted benzyl;


$R^1$  and  $R^2$  independently of one another

hydrogen, halogen,  $OR^{48}$ ,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_3$ - $C_{10}$ -alkynyl,  $C_3$ - $C_7$ -cycloalkyl, phenyl, benzyl or  $C_5$ - $C_7$ -cycloalkenyl, whereas each of the last-mentioned 7 groups can be substituted with any combination of one to six halogen atoms, one to three  $C_1$ - $C_6$ -alkoxy groups, one or two  $C_1$ - $C_8$ -haloalkoxy groups, one or two cyano groups, one or two  $C_3$ - $C_7$ -cycloalkyl groups, one or two  $C(O)R^{49}$  groups, one or two  $CO-OR^{50}$  groups, one or two  $CO-SR^{51}$  groups, one or two  $CO-NR^{52}R^{53}$  groups, one to three  $OR^{54}$  groups, one to three  $SR^{54}$  groups, one optionally substituted four to 10-membered monocyclic or fused bicyclic heterocyclic ring, one or two optionally substituted phenyl groups or one or two optionally substituted benzyl groups,


or  $R^1$  and  $R^2$  together with the atom to which they are attached form a 3- to 7-membered heterocyclic ring;

Q is selected from









wherein

A<sup>1</sup> to A<sup>17</sup> are each independently oxygen or sulfur;

30 R<sup>3</sup>, R<sup>4</sup>, R<sup>7</sup>, R<sup>8</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>18</sup>, R<sup>19</sup>, R<sup>27</sup>, R<sup>29</sup>, R<sup>32</sup>, R<sup>33</sup>, R<sup>38</sup>, R<sup>39</sup>, R<sup>44</sup>, R<sup>45</sup>, R<sup>46</sup> and R<sup>47</sup> are each independently

hydrogen, cyano, amino, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, benzyl, OR<sup>55</sup>, C<sub>1</sub>-C<sub>3</sub>-cyanoalkyl, or

35 R<sup>3</sup> and R<sup>4</sup>, R<sup>7</sup> and R<sup>8</sup>, R<sup>11</sup> and R<sup>12</sup>, R<sup>18</sup> and R<sup>19</sup> or R<sup>46</sup> and R<sup>47</sup> may be taken together with the atoms to which they are attached to represent a four- to seven-membered ring, optionally interrupted by oxygen, sulfur or nitrogen and optionally substituted with one or more halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl groups;

40 R<sup>5</sup>, R<sup>6</sup>, R<sup>9</sup>, R<sup>10</sup>, R<sup>15</sup>, R<sup>16</sup>, R<sup>20</sup>, R<sup>21</sup>, R<sup>30</sup>, R<sup>31</sup>, R<sup>35</sup>, R<sup>36</sup>, R<sup>41</sup>, R<sup>42</sup> and R<sup>43</sup> are each independently hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, OR<sup>56</sup>, S(O)<sub>n</sub>R<sup>57</sup>, O-SO<sub>2</sub>-R<sup>57</sup>, NR<sup>58</sup>R<sup>59</sup> or

45

R<sup>5</sup> and R<sup>6</sup>, R<sup>9</sup> and R<sup>10</sup>, R<sup>15</sup> and R<sup>16</sup>, R<sup>20</sup> and R<sup>21</sup> or R<sup>30</sup> and R<sup>31</sup> may be taken together with the atoms to which they are attached to represent a four- to seven membered ring optionally substituted with one or more halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl groups;

5

R<sup>13</sup>, R<sup>14</sup>, R<sup>22</sup>, R<sup>23</sup>, R<sup>25</sup> and R<sup>26</sup> are each independently hydrogen, halogen or C<sub>1</sub>-C<sub>6</sub>-alkyl;

10 R<sup>17</sup>, R<sup>28</sup>, R<sup>34</sup>, R<sup>37</sup> or R<sup>40</sup> are each independently hydrogen, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, OR<sup>60</sup> or SR<sup>61</sup>;

R<sup>24</sup> is hydrogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>2</sub>-C<sub>4</sub>-alkenyl, C<sub>3</sub>-C<sub>4</sub>-alkynyl, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy or amino;

15

R<sup>48</sup>, R<sup>49</sup>, R<sup>50</sup>, R<sup>51</sup>, R<sup>52</sup>, R<sup>53</sup>, R<sup>54</sup>, R<sup>55</sup>, R<sup>56</sup>, R<sup>57</sup>, R<sup>58</sup>, R<sup>59</sup>, R<sup>60</sup> and R<sup>61</sup> are independently of one another hydrogen, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, optionally substituted phenyl or

20

optionally substituted benzyl;


n is zero, 1 or 2;

25 and the agriculturally useful salts of the compounds I.

Furthermore, the invention relates to

- the use of the compounds I as herbicides and/or for the desiccation/defoliation of plants,
- 30 - herbicidal compositions and compositions for the desiccation/defoliation of plants which comprise compounds I as active substances,
- processes for preparing the compounds I and herbicidal compositions and compositions for the desiccation/defoliation of plants using the compounds I,
- 35 - methods for controlling undesirable vegetation and for the desiccation/defoliation of plants using the compounds I, and
- novel intermediates of the formula II.

40



(II)

wherein Q, X<sup>1</sup> and X<sup>2</sup> are as defined hereinabove, with the proviso that Q must be other than Q<sup>21</sup>.

45

Preferred compounds of the formulae I and II can be seen from the sub-claims and from the description which follows.

- Depending on the substitution pattern, the compounds of the  
5 formula I can contain one or more chiral centers, in which case they exist in the form of enantiomer or diastereomer mixtures. This invention provides both the pure enantiomers or diastereomers and mixtures thereof.
- 10 Agriculturally useful salts are in particular the salts of those cations and the acid addition salts of those acids whose cations and anions, respectively, do not adversely affect the herbicidal activity of the compounds I. Suitable cations are therefore in particular the ions of the alkali metals, preferably sodium and  
15 potassium, of the alkaline earth metals, preferably calcium, magnesium and barium, and of the transition metals, preferably manganese, copper, zinc and iron, and the ammonium ion, which may carry one to four C<sub>1</sub>-C<sub>4</sub>-alkyl substituents, and/or one phenyl or benzyl substituent, preferably diisopropylammonium,  
20 tetramethylammonium, tetrabutylammonium, trimethylbenzylammonium, and furthermore phosphonium ions, sulfonium ions, preferably tri(C<sub>1</sub>-C<sub>4</sub>-alkyl)sulfonium and sulfoxonium ions, preferably tri(C<sub>1</sub>-C<sub>4</sub>-alkyl)sulfoxonium.
- 25 Anions of useful acid addition salts are primarily chloride, bromide, fluoride, hydrogensulfate, sulfate, dihydrogenphosphate, hydrogenphosphate, phosphate, nitrate, hydrogencarbonate, carbonate, hexafluorosilicate, hexafluorophosphate, benzoate, and the anions of C<sub>1</sub>-C<sub>4</sub>-alkanoic acids, preferably formate, acetate,  
30 propionate and butyrate.

The organic moieties mentioned for the substituents X<sup>3</sup> and R<sup>1</sup> to R<sup>61</sup> or as radicals on phenyl or heterocyclic rings are collective terms for individual enumerations of each of the group members,  
35 as is the meaning halogen. All carbon chains, ie. all alkyl, haloalkyl, alkenyl, alkynyl and phenylalkyl moieties can be straight-chain or branched.

The terms haloalkyl, haloalkoxy and haloalkenyl as used in the  
40 specification and claims designate an alkyl group, an alkoxy group or an alkenyl group substituted with one or more halogen atoms, respectively. The halogen atoms may be the same or different.

Halogenated substituents preferably have attached to them one to  
45 five identical or different halogen atoms.

In formula I above, 4- to 10-membered monocyclic or fused bicyclic, heterocyclic rings include, but are not limited to, benzimidazole, imidazole, imidazoline-2-thione, indole, isatoic anhydride, morpholine, piperazine, piperidine, purine, pyrazole, pyrrole, pyrrolidine and 1,2,4-triazole rings, wherein each ring is optionally substituted with one or more groups independently selected from halogen, cyano, nitro, amino, hydroxyl, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl, C<sub>1</sub>-C<sub>4</sub>-alkoxy, C<sub>1</sub>-C<sub>4</sub>-haloalkoxy or C<sub>1</sub>-C<sub>4</sub>-haloalkyl-sulfonyl groups.

10

When the terms phenyl or benzyl are designated as being optionally substituted, the substituents which are optionally present may be any one or more of those customarily employed in the development of pesticidal compounds and/or the modification of such compounds to influence their structure/activity, persistence, penetration or other property. Specific examples of such substituents include, for example, halogen atoms, nitro, cyano, thiocyanato, cyanato, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, formyl, alkoxy-carbonyl, carboxyl, alkanoyl, alkylthio, alkylsulphinyl, alkyl-sulfonyl, carbamoyl, alkylamido, phenyl, phenoxy, benzyl, benzyl-oxy, heterocyclyl, especially furyl, and cycloalkyl, especially cyclopropyl, groups. Typically, zero to three substituents may be present. When any of the foregoing substituents represents or contains an alkyl substituent group, this may be linear or branched and may contain up to 12, preferably up to 6, and especially up to 4 carbon atoms.

In formula I above, 3- to 7-membered heterocyclic rings include, but are not limited to, imidazole and phthalimide rings wherein each ring is optionally substituted with any combination of one to three halogen atoms, one to three C<sub>1</sub>-C<sub>4</sub>-alkyl groups, one to three C<sub>1</sub>-C<sub>4</sub>-haloalkyl groups, one to three C<sub>1</sub>-C<sub>4</sub>-alkoxy groups, or one to three C<sub>1</sub>-C<sub>4</sub>-haloalkoxy groups.

35

The uracil substituted phenyl sulfamoyl carboxamides I possess an unexpected level of herbicidal activity and surprising crop selectivity.

40 Examples of individual meanings are:

- halogen: fluorine, chlorine, bromine or iodine, preferably fluorine or chlorine;
- 45 - C<sub>1</sub>-C<sub>4</sub>-alkyl; CH<sub>3</sub>, C<sub>2</sub>H<sub>5</sub>, CH<sub>2</sub>-C<sub>2</sub>H<sub>5</sub>, CH(CH<sub>3</sub>)<sub>2</sub>, n-C<sub>4</sub>H<sub>9</sub>, CH(CH<sub>3</sub>)-C<sub>2</sub>H<sub>5</sub>, CH<sub>2</sub>-CH(CH<sub>3</sub>)<sub>2</sub> or C(CH<sub>3</sub>)<sub>3</sub>;

- C<sub>1</sub>-C<sub>6</sub>-alkyl and the alkyl moiety of C<sub>1</sub>-C<sub>6</sub>-alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl: methyl, ethyl, n-propyl, 1-methylethyl, n-butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl or 1-ethyl-2-methylpropyl, preferably C<sub>1</sub>-C<sub>4</sub>-alkyl, in particular methyl or ethyl;
- 15 - C<sub>1</sub>-C<sub>3</sub>-cyanoalkyl: CH<sub>2</sub>CN, 1-cyanoethyl, 2-cyanoethyl, 1-cyanoprop-1-yl, 2-cyanoprop-1-yl, 3-cyanoprop-1-yl or 1-(CH<sub>2</sub>CN)eth-1-yl;
- 20 - C<sub>1</sub>-C<sub>6</sub>-haloalkyl: C<sub>1</sub>-C<sub>6</sub>-alkyl as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, eg. CH<sub>2</sub>F, CHF<sub>2</sub>, CF<sub>3</sub>, CH<sub>2</sub>Cl, CH(Cl)<sub>2</sub>, C(Cl)<sub>3</sub>, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, C<sub>2</sub>F<sub>5</sub>, 3-fluoropropyl, 3-chloropropyl or fluoromethyl or 1,2-dichloroethyl;
- 30
- C<sub>2</sub>-C<sub>6</sub>-alkenyl: ethenyl, prop-1-en-1-yl, prop-2-en-1-yl, 1-methylethenyl, n-buten-1-yl, n-buten-2-yl, n-buten-3-yl, 1-methylprop-1-en-1-yl, 2-methylprop-1-en-1-yl, 1-methylprop-2-en-1-yl, 2-methylprop-2-en-1-yl, n-penten-1-yl, n-penten-2-yl, n-penten-3-yl, n-penten-4-yl, 1-methylbut-1-en-1-yl, 2-methylbut-1-en-1-yl, 3-methylbut-1-en-1-yl, 1-methylbut-2-en-1-yl, 2-methylbut-2-en-1-yl, 3-methylbut-2-en-1-yl, 1-methylbut-3-en-1-yl, 2-methylbut-3-en-1-yl, 3-methylbut-3-en-1-yl, 1,1-dimethylprop-2-en-1-yl, 1,2-dimethylprop-1-en-1-yl, 1,2-dimethylprop-2-en-1-yl, 1-ethylprop-1-en-2-yl, 1-ethylprop-2-en-1-yl, n-hex-1-en-1-yl, n-hex-2-en-1-yl, n-hex-3-en-1-yl, n-hex-4-en-1-yl, n-hex-5-en-1-yl, 1-methylpent-1-en-1-yl, 2-methylpent-1-en-1-yl, 3-methylpent-1-en-1-yl, 4-methylpent-1-en-1-yl, 1-methylpent-2-en-1-yl,
- 35
- 40
- 45

- 2-methylpent-2-en-1-yl, 3-methylpent-2-en-1-yl,  
 4-methylpent-2-en-1-yl, 1-methylpent-3-en-1-yl,  
 2-methylpent-3-en-1-yl, 3-methylpent-3-en-1-yl,  
 4-methylpent-3-en-1-yl, 1-methylpent-4-en-1-yl,  
 5 2-methylpent-4-en-1-yl, 3-methylpent-4-en-1-yl,  
 4-methylpent-4-en-1-yl, 1,1-dimethylbut-2-en-1-yl,  
 1,1-dimethylbut-3-en-1-yl, 1,2-dimethylbut-1-en-1-yl,  
 1,2-dimethylbut-2-en-1-yl, 1,2-dimethylbut-3-en-1-yl,  
 1,3-dimethylbut-1-en-1-yl, 1,3-dimethylbut-2-en-1-yl,  
 10 1,3-dimethylbut-3-en-1-yl, 2,2-dimethylbut-3-en-1-yl,  
 2,3-dimethylbut-1-en-1-yl, 2,3-dimethylbut-2-en-1-yl,  
 2,3-dimethylbut-3-en-1-yl, 3,3-dimethylbut-1-en-1-yl,  
 3,3-dimethylbut-2-en-1-yl, 1-ethylbut-1-en-1-yl,  
 1-ethylbut-2-en-1-yl, 1-ethylbut-3-en-1-yl,  
 15 2-ethylbut-1-en-1-yl, 2-ethylbut-2-en-1-yl,  
 2-ethylbut-3-en-1-yl, 1,1,2-trimethylprop-2-en-1-yl,  
 1-ethyl-1-methylprop-2-en-1-yl,  
 1-ethyl-2-methylprop-1-en-1-yl or  
 1-ethyl-2-methylprop-2-en-1-yl, preferably C<sub>3</sub>- or C<sub>4</sub>-alkenyl;  
 20
- C<sub>2</sub>-C<sub>6</sub>-haloalkenyl: C<sub>2</sub>-C<sub>6</sub>-alkenyl as mentioned above which is partially or fully substituted by fluorine, chlorine, bromine and/or iodine, ie. for example 2-chloroallyl, 3-chloroallyl,  
 25 2,3-dichloroallyl, 3,3-dichloroallyl, 2,3,4-trichloroallyl,  
 2,3-dichlorobut-2-enyl, 2-bromoallyl, 3-bromoallyl,  
 2,3-dibromoallyl, 3,3-dibromoallyl, 2,3,3-tribromoallyl or  
 2,3-dibromobut-2-enyl;
- 30 - C<sub>3</sub>-C<sub>6</sub>-alkynyl: prop-1-yn-1-yl, prop-2-yn-3-yl,  
 n-but-1-yn-1-yl, n-but-1-yn-4-yl, n-but-2-yn-1-yl,  
 n-pent-1-yn-1-yl, n-pent-1-yn-3-yl, n-pent-1-yn-4-yl,  
 n-pent-1-yn-5-yl, n-pent-2-yn-1-yl, n-pent-2-yn-4-yl,  
 n-pent-2-yn-5-yl, 3-methylbut-1-yn-1-yl,  
 35 3-methylbut-1-yn-3-yl, 3-methylbut-1-yn-4-yl,  
 n-hex-1-yn-1-yl, n-hex-1-yn-3-yl, n-hex-1-yn-4-yl,  
 n-hex-1-yn-5-yl, n-hex-1-yn-6-yl, n-hex-2-yn-1-yl,  
 n-hex-2-yn-4-yl, n-hex-2-yn-5-yl, n-hex-2-yn-6-yl,  
 n-hex-3-yn-1-yl, n-hex-3-yn-2-yl, 3-methylpent-1-yn-1-yl,  
 40 3-methylpent-1-yn-3-yl, 3-methylpent-1-yn-4-yl,  
 3-methylpent-1-yn-5-yl, 4-methylpent-1-yn-1-yl,  
 4-methylpent-2-yn-4-yl or 4-methylpent-2-yn-5-yl, preferably  
 C<sub>3</sub>- or C<sub>4</sub>-alkynyl, in particular prop-2-yn-3-yl;
- 45 - phenyl-C<sub>1</sub>-C<sub>6</sub>-alkyl: for example benzyl, 1-phenyleth-1-yl,  
 2-phenyleth-1-yl, 1-phenylprop-1-yl, 2-phenylprop-1-yl,  
 3-phenylprop-1-yl, 1-phenylprop-2-yl, 2-phenylprop-2-yl,

- 1-phenylbut-1-yl, 2-phenylbut-1-yl, 3-phenylbut-1-yl,  
4-phenylbut-1-yl, 1-phenylbut-2-yl, 2-phenylbut-2-yl,  
1-phenylbut-3-yl, 2-phenylbut-3-yl,  
1-phenyl-2-methylprop-3-yl, 2-phenyl-2-methylprop-3-yl,  
5 3-phenyl-2-methylprop-3-yl or 2-benzylprop-2-yl, preferably  
phenyl-C<sub>1</sub>-C<sub>4</sub>-alkyl, in particular 2-phenyleth-1-yl;
- C<sub>1</sub>-C<sub>6</sub>-alkoxy and the alkoxy moiety of  
C<sub>1</sub>-C<sub>6</sub>-Alkoxy-C<sub>1</sub>-C<sub>6</sub>-alkyl: methoxy, ethoxy, n-propoxy,  
10 1-methylethoxy, n-butoxy, 1-methylpropoxy, 2-methylpropoxy,  
1,1-dimethylethoxy, n-pentoxy, 1-methylbutoxy,  
2-methylbutoxy, 3-methylbutoxy, 1,1-dimethylpropoxy,  
1,2-dimethylpropoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy,  
n-hexoxy, 1-methylpentoxy, 2-methylpentoxy, 3-methylpentoxy,  
15 4-methylpentoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy,  
1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy,  
3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy,  
1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy,  
1-ethyl-1-methylpropoxy or 1-ethyl-2-methylpropoxy,  
20 preferably C<sub>1</sub>-C<sub>4</sub>-alkoxy, in particular OCH<sub>3</sub>, OC<sub>2</sub>H<sub>5</sub> or  
OCH(CH<sub>3</sub>)<sub>2</sub>;
- C<sub>3</sub>-C<sub>7</sub>-cycloalkyl: cyclopropyl, cyclobutyl, cyclopentyl,  
25 cyclohexyl and cycloheptyl;
- C<sub>5</sub>-C<sub>7</sub>-Cycloalkenyl: cyclopent-1-enyl, cyclopent-2-enyl,  
cyclopent-3-enyl, cyclohex-1-enyl, cyclohex-2-enyl,  
cyclohex-3-enyl, cyclohept-1-enyl, cyclohept-2-enyl,  
30 cyclohept-3-enyl, cyclohept-4-enyl, cyclooct-1-enyl,  
cyclooct-2-enyl, cyclooct-3-enyl and cyclooct-4-enyl.
- 3- to 7-membered heterocycle is a saturated, partially or fully  
unsaturated or aromatic heterocycle having one to three hetero  
35 atoms selected from a group consisting of
- one to three nitrogens,
  - one or two oxygens and
  - one or two sulfur atoms.
- 40 With a view to the use of the compounds I as herbicides and/or  
compounds which have a desiccant/defoliant action, the variables  
preferably have the following meanings, to be precise in each  
case alone or in combination:
- 45 X<sup>1</sup> is hydrogen or halogen, in particular hydrogen or chlorine;

- $x^2$  is cyano or halogen, in particular cyano or chlorine;
- $x^3$  is hydrogen;
- <sup>5</sup>  $Q$  is  $Q^5$ ,  $Q^7$ ,  $Q^{21}$ ,  $Q^{22}$ ,  $Q^{27}$ ,  $Q^{32}$ ,  $Q^{38}$ ,  $Q^{39}$  or  $Q^{40}$ ;
- $A^1$  is oxygen;
- <sup>10</sup>  $A^3$ ,  $A^4$  are, independently of one another, oxygen;
- $A^8$ ,  $A^9$  are, independently of one another, oxygen;
- <sup>15</sup>  $A^{10}$ ,  $A^{11}$  are, independently of one another, oxygen;
- $A^{12}$  is sulfur;
- $A^{13}$  is oxygen;
- <sup>20</sup>  $A^{15}$  is sulfur;
- $R^1$  is  $C_1$ - $C_4$ -alkyl;
- <sup>25</sup>  $R^2$  is  $C_1$ - $C_6$ -alkyl,  $C_2$ - $C_6$ -alkenyl or  $C_2$ - $C_6$ -alkynyl;
- $R^7$  is amino,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkoxy or  $C_1$ - $C_6$ -haloalkoxy;
- <sup>30</sup>  $R^8$  is  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_3$ - $C_7$ -cycloalkyl,  $C_2$ - $C_6$ -alkenyl or  $C_1$ - $C_6$ -haloalkoxy;
- <sup>35</sup>  $R^7$  and  $R^8$  may be taken together with the atoms to which they are attached to represent a four to seven membered ring, optionally interrupted by oxygen, sulfur or nitrogen;
- $R^9$ ,  $R^{10}$  are, independently of one another, hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -alkoxy or together with the atoms to which they are attached to represent a 5- or 6-membered ring;
- <sup>40</sup>  $R^{29}$  is hydrogen, amino or  $C_1$ - $C_6$ -alkyl;
- <sup>45</sup>  $R^{30}$  is  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -haloalkoxy or  $C_1$ - $C_6$ -alkylsulfonyl and



## 12

R<sup>31</sup> is hydrogen, amino, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl or C<sub>2</sub>-C<sub>6</sub>-alkenyl or

R<sup>30</sup> and R<sup>31</sup> together with the atoms to which they are attached to represent a 5- or 6-membered ring;

5

R<sup>32</sup> is hydrogen, amino, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>2</sub>-C<sub>6</sub>-alkenyl;

10 R<sup>33</sup> is hydrogen, amino, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl or C<sub>2</sub>-C<sub>6</sub>-alkenyl;

R<sup>34</sup> is hydrogen oder C<sub>1</sub>-C<sub>6</sub>-alkyl;

15 R<sup>35</sup> is C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy or C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl;

R<sup>36</sup> is hydrogen, amino, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl or C<sub>2</sub>-C<sub>6</sub>-alkenyl;

20

R<sup>37</sup> is hydrogen, cyano, halogen, C<sub>1</sub>-C<sub>6</sub>-alkyl or C<sub>1</sub>-C<sub>6</sub>-alkoxy;

R<sup>38</sup> is cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy or C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl;

25

R<sup>39</sup> is cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy or C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl;

30 R<sup>40</sup> is halogen;

R<sup>41</sup> is hydrogen, amino or C<sub>1</sub>-C<sub>6</sub>-alkyl;

35 R<sup>42</sup> is C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkoxy, C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyl or C<sub>1</sub>-C<sub>6</sub>-alkylsulfonyloxy;

R<sup>43</sup> is hydrogen, amino or C<sub>1</sub>-C<sub>6</sub>-alkyl;

40 R<sup>44</sup> is hydrogen, amino or C<sub>1</sub>-C<sub>6</sub>-alkyl;

R<sup>45</sup> is hydrogen, amino or C<sub>1</sub>-C<sub>6</sub>-alkyl;


45

$R^{46}$ ,  $R^{47}$  are, independently of one another,  $C_1$ - $C_6$ -haloalkyl or together with the nitrogen atoms to which they are attached to represent a 5- or 6-membered ring, optionally interrupted by one oxygen or sulfur ring member.

5

Very especially preferred are the compounds of the formula Ia ( $\square$  I where  $X^1$  = fluorine;  $X^2$  = chlorine;  $Q = Q^{21}$ ;  $X^3$  = hydrogen;  $A$ ,  $A^8$ ,  $A^9$  = oxygen;  $R^{29}$  = methyl;  $R^{30}$  = trifluoromethyl;  $R^{31}$  = hydrogen)

10



Ia,

15

in particular the compounds of Table 1:

Table 1

20

No.	$R^1$	$R^2$
Ia.1	H	$CH_3$
Ia.2	H	$C_2H_5$
Ia.3	H	$CH_2CH_2-Cl$
25 Ia.4	H	$CH_2CH_2-CN$
Ia.5	H	$CH_2-CO-OCH_3$
Ia.6	H	$CH_2-CO-OC_2H_5$
Ia.7	H	$CH(CH_3)-CO-OCH_3$
30 Ia.8	H	$CH_2CH_2-OCH_3$
Ia.9	H	$CH_2-C_2H_5$
Ia.10	H	$CH_2CH_2-C_2H_5$
Ia.11	H	$CH(CH_3)_2$
35 Ia.12	H	$CH(CH_3)-C_2H_5$
Ia.13	H	$CH_2-CH(CH_3)_2$
Ia.14	H	$C(CH_3)_3$
Ia.15	H	$CH(CH_3)-CH_2-C_2H_5$
Ia.16	H	$CH_2-CH(CH_3)-C_2H_5$
40 Ia.17	H	$CH_2CH_2-CH(CH_3)_2$
Ia.18	H	$CH_2-CH=CH_2$
Ia.19	H	$CH(CH_3)=CH_2$
Ia.20	H	$CH_2=CH-CH_3$
45 Ia.21	H	$CH_2-C\square CH$
Ia.22	H	$CH(CH_3)-C\square CH$

No.	R <sup>1</sup>	R <sup>2</sup>
Ia.23	H	Cyclopropyl
Ia.24	H	CH <sub>2</sub> -Cyclopropyl
5 Ia.25	H	Cyclopentyl
Ia.26	H	CH <sub>2</sub> -Cyclopentyl
Ia.27	H	CH <sub>2</sub> -(1,3-Dioxolanyl)
Ia.28	H	CH <sub>2</sub> -(2-Furyl)
10 Ia.29	H	CH <sub>2</sub> -(3-Furyl)
Ia.30	H	CH <sub>2</sub> -(2-Thienyl)
Ia.31	H	CH <sub>2</sub> -(3-Thienyl)
Ia.32	H	Phenyl
Ia.33	H	2-Chlorophenyl
15 Ia.34	H	3-Chlorophenyl
Ia.35	H	4-Chlorophenyl
Ia.36	H	2-Fluorophenyl
Ia.37	H	3-Fluorophenyl
20 Ia.38	H	4-Fluorophenyl
Ia.39	H	2-Methylphenyl
Ia.40	H	3-Methylphenyl
Ia.41	H	4-Methylphenyl
25 Ia.42	H	2-Methoxyphenyl
Ia.43	H	3-Methoxyphenyl
Ia.44	H	4-Methoxyphenyl
Ia.45	H	2-(Methoxycarbonyl)phenyl
Ia.46	H	3-(Methoxycarbonyl)phenyl
30 Ia.47	H	4-(Methoxycarbonyl)phenyl
Ia.48	H	2-Nitrophenyl
Ia.49	H	3-Nitrophenyl
Ia.50	H	4-Nitrophenyl
35 Ia.51	H	2-(Dimethylamino)phenyl
Ia.52	H	3-(Dimethylamino)phenyl
Ia.53	H	4-(Dimethylamino)phenyl
Ia.54	H	2-(Trifluoromethyl)phenyl
40 Ia.55	H	3-(Trifluoromethyl)phenyl
Ia.56	H	4-(Trifluoromethyl)phenyl
Ia.57	H	3-(Phenoxy)phenyl
Ia.58	H	4-(Phenoxy)phenyl
45 Ia.59	H	2,4-Difluorophenyl
Ia.60	H	2,4-Dichlorophenyl
Ia.61	H	3,4-Difluorophenyl

	No.	R <sup>1</sup>	R <sup>2</sup>
5	Ia.62	H	3,4-Dichlorophenyl
	Ia.63	H	3,5-Difluorophenyl
	Ia.64	H	3,5-Dichlorophenyl
	Ia.65	H	2-Pyridyl
	Ia.66	H	3-Pyridyl
10	Ia.67	H	4-Pyridyl
	Ia.68	H	$\alpha$ -Naphthyl
	Ia.69	H	Benzyl
	Ia.70	H	2-Chlorobenzyl
	Ia.71	H	3-Chlorobenzyl
15	Ia.72	H	4-Chlorobenzyl
	Ia.73	H	2-Methoxybenzyl
	Ia.74	H	3-Methoxybenzyl
	Ia.75	H	4-Methoxybenzyl
	Ia.76	CH <sub>3</sub>	CH <sub>3</sub>
20	Ia.77	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>
	Ia.78	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> -Cl
	Ia.79	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> -CN
	Ia.80	CH <sub>3</sub>	CH <sub>2</sub> -CO-OCH <sub>3</sub>
	Ia.81	CH <sub>3</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>
25	Ia.82	CH <sub>3</sub>	CH(CH <sub>3</sub> )-CO-OCH <sub>3</sub>
	Ia.83	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> -OCH <sub>3</sub>
	Ia.84	CH <sub>3</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.85	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.86	CH <sub>3</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>
30	Ia.87	CH <sub>3</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.88	CH <sub>3</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.89	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
	Ia.90	CH <sub>3</sub>	CH(CH <sub>3</sub> )-CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.91	CH <sub>3</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
35	Ia.92	CH <sub>3</sub>	CH <sub>2</sub> CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.93	CH <sub>3</sub>	CH <sub>2</sub> -CH=CH <sub>2</sub>
	Ia.94	CH <sub>3</sub>	CH(CH <sub>3</sub> )=CH <sub>2</sub>
	Ia.95	CH <sub>3</sub>	CH <sub>2</sub> =CH-CH <sub>3</sub>
	Ia.96	CH <sub>3</sub>	CH <sub>2</sub> -C $\square$ CH
40	Ia.97	CH <sub>3</sub>	CH(CH <sub>3</sub> )-C $\square$ CH
	Ia.98	CH <sub>3</sub>	Cyclopropyl
	Ia.99	CH <sub>3</sub>	CH <sub>2</sub> -Cyclopropyl
	Ia.100	CH <sub>3</sub>	Cyclopentyl

No.	R <sup>1</sup>	R <sup>2</sup>
Ia.101	CH <sub>3</sub>	CH <sub>2</sub> -Cyclopentyl
Ia.102	CH <sub>3</sub>	CH <sub>2</sub> -(1,3-Dioxolanyl)
5 Ia.103	CH <sub>3</sub>	CH <sub>2</sub> -(2-Furyl)
Ia.104	CH <sub>3</sub>	CH <sub>2</sub> -(3-Furyl)
Ia.105	CH <sub>3</sub>	CH <sub>2</sub> -(2-Thienyl)
Ia.106	CH <sub>3</sub>	CH <sub>2</sub> -(3-Thienyl)
10 Ia.107	CH <sub>3</sub>	Phenyl
Ia.108	CH <sub>3</sub>	2-Chlorophenyl
Ia.109	CH <sub>3</sub>	3-Chlorophenyl
Ia.110	CH <sub>3</sub>	4-Chlorophenyl
Ia.111	CH <sub>3</sub>	2-Fluorophenyl
15 Ia.112	CH <sub>3</sub>	3-Fluorophenyl
Ia.113	CH <sub>3</sub>	4-Fluorophenyl
Ia.114	CH <sub>3</sub>	2-Methylphenyl
Ia.115	CH <sub>3</sub>	3-Methylphenyl
20 Ia.116	CH <sub>3</sub>	4-Methylphenyl
Ia.117	CH <sub>3</sub>	2-Methoxyphenyl
Ia.118	CH <sub>3</sub>	3-Methoxyphenyl
Ia.119	CH <sub>3</sub>	4-Methoxyphenyl
25 Ia.120	CH <sub>3</sub>	2-(Methoxycarbonyl)phenyl
Ia.121	CH <sub>3</sub>	3-(Methoxycarbonyl)phenyl
Ia.122	CH <sub>3</sub>	4-(Methoxycarbonyl)phenyl
Ia.123	CH <sub>3</sub>	2-Nitrophenyl
Ia.124	CH <sub>3</sub>	3-Nitrophenyl
30 Ia.125	CH <sub>3</sub>	4-Nitrophenyl
Ia.126	CH <sub>3</sub>	2-(Dimethylamino)phenyl
Ia.127	CH <sub>3</sub>	3-(Dimethylamino)phenyl
Ia.128	CH <sub>3</sub>	4-(Dimethylamino)phenyl
35 Ia.129	CH <sub>3</sub>	2-(Trifluoromethyl)phenyl
Ia.130	CH <sub>3</sub>	3-(Trifluoromethyl)phenyl
Ia.131	CH <sub>3</sub>	4-(Trifluoromethyl)phenyl
Ia.132	CH <sub>3</sub>	3-(Phenoxy)phenyl
40 Ia.133	CH <sub>3</sub>	4-(Phenoxy)phenyl
Ia.134	CH <sub>3</sub>	2,4-Difluorophenyl
Ia.135	CH <sub>3</sub>	2,4-Dichlorophenyl
Ia.136	CH <sub>3</sub>	3,4-Difluorophenyl
Ia.137	CH <sub>3</sub>	3,4-Dichlorophenyl
45 Ia.138	CH <sub>3</sub>	3,5-Difluorophenyl
Ia.139	CH <sub>3</sub>	3,5-Dichlorophenyl

	No.	R <sup>1</sup>	R <sup>2</sup>
5	Ia.140	CH <sub>3</sub>	2-Pyridyl
	Ia.141	CH <sub>3</sub>	3-Pyridyl
	Ia.142	CH <sub>3</sub>	4-Pyridyl
	Ia.143	CH <sub>3</sub>	$\alpha$ -Naphthyl
	Ia.144	CH <sub>3</sub>	Benzyl
10	Ia.145	CH <sub>3</sub>	2-Chlorobenzyl
	Ia.146	CH <sub>3</sub>	3-Chlorobenzyl
	Ia.147	CH <sub>3</sub>	4-Chlorobenzyl
	Ia.148	CH <sub>3</sub>	2-Methoxybenzyl
	Ia.149	CH <sub>3</sub>	3-Methoxybenzyl
15	Ia.150	CH <sub>3</sub>	4-Methoxybenzyl
	Ia.151	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>
	Ia.152	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -Cl
	Ia.153	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -CN
	Ia.154	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CO-OCH <sub>3</sub>
20	Ia.155	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>
	Ia.156	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-CO-OCH <sub>3</sub>
	Ia.157	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -OCH <sub>3</sub>
	Ia.158	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.159	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
25	Ia.160	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.161	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.162	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.163	C <sub>2</sub> H <sub>5</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
	Ia.164	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
30	Ia.165	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.166	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.167	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH=CH <sub>2</sub>
	Ia.168	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )=CH <sub>2</sub>
	Ia.169	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> =CH-CH <sub>3</sub>
35	Ia.170	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C $\square$ CH
	Ia.171	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-C $\square$ CH
	Ia.172	C <sub>2</sub> H <sub>5</sub>	Cyclopropyl
	Ia.173	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -Cyclopropyl
	Ia.174	C <sub>2</sub> H <sub>5</sub>	Cyclopentyl
40	Ia.175	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -Cyclopentyl
	Ia.176	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(1,3-Dioxolanyl)
	Ia.177	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(2-Furyl)
	Ia.178	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(3-Furyl)

No.	R <sup>1</sup>	R <sup>2</sup>
Ia.179	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(2-Thienyl)
Ia.180	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(3-Thienyl)
5 Ia.181	C <sub>2</sub> H <sub>5</sub>	Phenyl
Ia.182	C <sub>2</sub> H <sub>5</sub>	2-Chlorophenyl
Ia.183	C <sub>2</sub> H <sub>5</sub>	3-Chlorophenyl
Ia.184	C <sub>2</sub> H <sub>5</sub>	4-Chlorophenyl
10 Ia.185	C <sub>2</sub> H <sub>5</sub>	2-Fluorophenyl
Ia.186	C <sub>2</sub> H <sub>5</sub>	3-Fluorophenyl
Ia.187	C <sub>2</sub> H <sub>5</sub>	4-Fluorophenyl
Ia.188	C <sub>2</sub> H <sub>5</sub>	2-Methylphenyl
Ia.189	C <sub>2</sub> H <sub>5</sub>	3-Methylphenyl
15 Ia.190	C <sub>2</sub> H <sub>5</sub>	4-Methylphenyl
Ia.191	C <sub>2</sub> H <sub>5</sub>	2-Methoxyphenyl
Ia.192	C <sub>2</sub> H <sub>5</sub>	3-Methoxyphenyl
Ia.193	C <sub>2</sub> H <sub>5</sub>	4-Methoxyphenyl
20 Ia.194	C <sub>2</sub> H <sub>5</sub>	2-(Methoxycarbonyl)phenyl
Ia.195	C <sub>2</sub> H <sub>5</sub>	3-(Methoxycarbonyl)phenyl
Ia.196	C <sub>2</sub> H <sub>5</sub>	4-(Methoxycarbonyl)phenyl
Ia.197	C <sub>2</sub> H <sub>5</sub>	2-Nitrophenyl
25 Ia.198	C <sub>2</sub> H <sub>5</sub>	3-Nitrophenyl
Ia.199	C <sub>2</sub> H <sub>5</sub>	4-Nitrophenyl
Ia.200	C <sub>2</sub> H <sub>5</sub>	2-(Dimethylamino)phenyl
Ia.201	C <sub>2</sub> H <sub>5</sub>	3-(Dimethylamino)phenyl
Ia.202	C <sub>2</sub> H <sub>5</sub>	4-(Dimethylamino)phenyl
30 Ia.203	C <sub>2</sub> H <sub>5</sub>	2-(Trifluoromethyl)phenyl
Ia.204	C <sub>2</sub> H <sub>5</sub>	3-(Trifluoromethyl)phenyl
Ia.205	C <sub>2</sub> H <sub>5</sub>	4-(Trifluoromethyl)phenyl
Ia.206	C <sub>2</sub> H <sub>5</sub>	3-(Phenoxy)phenyl
35 Ia.207	C <sub>2</sub> H <sub>5</sub>	4-(Phenoxy)phenyl
Ia.208	C <sub>2</sub> H <sub>5</sub>	2,4-Difluorophenyl
Ia.209	C <sub>2</sub> H <sub>5</sub>	2,4-Dichlorophenyl
Ia.210	C <sub>2</sub> H <sub>5</sub>	3,4-Difluorophenyl
40 Ia.211	C <sub>2</sub> H <sub>5</sub>	3,4-Dichlorophenyl
Ia.212	C <sub>2</sub> H <sub>5</sub>	3,5-Difluorophenyl
Ia.213	C <sub>2</sub> H <sub>5</sub>	3,5-Dichlorophenyl
Ia.214	C <sub>2</sub> H <sub>5</sub>	2-Pyridyl
45 Ia.215	C <sub>2</sub> H <sub>5</sub>	3-Pyridyl
Ia.216	C <sub>2</sub> H <sub>5</sub>	4-Pyridyl
Ia.217	C <sub>2</sub> H <sub>5</sub>	α-Naphthyl

	No.	R <sup>1</sup>	R <sup>2</sup>
	Ia.218	C <sub>2</sub> H <sub>5</sub>	Benzyl
	Ia.219	C <sub>2</sub> H <sub>5</sub>	2-Chlorobenzyl
5	Ia.220	C <sub>2</sub> H <sub>5</sub>	3-Chlorobenzyl
	Ia.221	C <sub>2</sub> H <sub>5</sub>	4-Chlorobenzyl
	Ia.222	C <sub>2</sub> H <sub>5</sub>	2-Methoxybenzyl
	Ia.223	C <sub>2</sub> H <sub>5</sub>	3-Methoxybenzyl
10	Ia.224	C <sub>2</sub> H <sub>5</sub>	4-Methoxybenzyl
	Ia.225	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>
	Ia.226	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -Cl
	Ia.227	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -CN
	Ia.228	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CO-OCH <sub>3</sub>
15	Ia.229	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>
	Ia.230	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-CO-OCH <sub>3</sub>
	Ia.231	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -OCH <sub>3</sub>
	Ia.232	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
20	Ia.233	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.234	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.235	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.236	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
25	Ia.237	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
	Ia.238	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.239	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.240	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
30	Ia.241	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH=CH <sub>2</sub>
	Ia.242	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )=CH <sub>2</sub>
	Ia.243	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> =CH-CH <sub>3</sub>
	Ia.244	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C□CH
	Ia.245	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-C□CH
35	Ia.246	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Cyclopropyl
	Ia.247	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -Cyclopropyl
	Ia.248	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Cyclopentyl
	Ia.249	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -Cyclopentyl
40	Ia.250	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(1,3-Dioxolanyl)
	Ia.251	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(2-Furyl)
	Ia.252	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(3-Furyl)
	Ia.253	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(2-Thienyl)
45	Ia.254	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(3-Thienyl)
	Ia.255	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Phenyl
	Ia.256	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Chlorophenyl



No.	R <sup>1</sup>	R <sup>2</sup>
Ia.257	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Chlorophenyl
Ia.258	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Chlorophenyl
5 Ia.259	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Fluorophenyl
Ia.260	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Fluorophenyl
Ia.261	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Fluorophenyl
Ia.262	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Methylphenyl
10 Ia.263	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Methylphenyl
Ia.264	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Methylphenyl
Ia.265	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Methoxyphenyl
Ia.266	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Methoxyphenyl
Ia.267	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Methoxyphenyl
15 Ia.268	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-(Methoxycarbonyl)phenyl
Ia.269	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Methoxycarbonyl)phenyl
Ia.270	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Methoxycarbonyl)phenyl
Ia.271	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Nitrophenyl
20 Ia.272	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Nitrophenyl
Ia.273	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Nitrophenyl
Ia.274	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-(Dimethylamino)phenyl
Ia.275	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Dimethylamino)phenyl
25 Ia.276	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Dimethylamino)phenyl
Ia.277	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-(Trifluoromethyl)phenyl
Ia.278	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Trifluoromethyl)phenyl
Ia.279	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Trifluoromethyl)phenyl
Ia.280	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Phenoxy)phenyl
30 Ia.281	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Phenoxy)phenyl
Ia.282	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2,4-Difluorophenyl
Ia.283	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2,4-Dichlorophenyl
Ia.284	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,4-Difluorophenyl
35 Ia.285	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,4-Dichlorophenyl
Ia.286	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,5-Difluorophenyl
Ia.287	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,5-Dichlorophenyl
Ia.288	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Pyridyl
40 Ia.289	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Pyridyl
Ia.290	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Pyridyl
Ia.291	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	$\alpha$ -Naphthyl
Ia.292	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Benzyl
45 Ia.293	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Chlorobenzyl
Ia.294	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Chlorobenzyl
Ia.295	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Chlorobenzyl

	No.	R <sup>1</sup>	R <sup>2</sup>
	Ia. 296	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Methoxybenzyl
	Ia. 297	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Methoxybenzyl
5	Ia. 298	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Methoxybenzyl
	Ia. 299	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -Cl
	Ia. 300	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -CN
	Ia. 301	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CO-OCH <sub>3</sub>
10	Ia. 302	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>
	Ia. 303	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-CO-OCH <sub>3</sub>
	Ia. 304	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -OCH <sub>3</sub>
	Ia. 305	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia. 306	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>
15	Ia. 307	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia. 308	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia. 309	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
	Ia. 310	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
20	Ia. 311	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia. 312	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia. 313	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -CH=CH <sub>2</sub>
	Ia. 314	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )=CH <sub>2</sub>
25	Ia. 315	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> =CH-CH <sub>3</sub>
	Ia. 316	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C $\square$ CH
	Ia. 317	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> )-C $\square$ CH
	Ia. 318	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Cyclopropyl
30	Ia. 319	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -Cyclopropyl
	Ia. 320	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Cyclopentyl
	Ia. 321	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -Cyclopentyl
	Ia. 322	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(1,3-Dioxolanyl)
	Ia. 323	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(2-Furyl)
35	Ia. 324	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(3-Furyl)
	Ia. 325	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(2-Thienyl)
	Ia. 326	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -(3-Thienyl)
	Ia. 327	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Phenyl
40	Ia. 328	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Chlorophenyl
	Ia. 329	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Chlorophenyl
	Ia. 330	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Chlorophenyl
	Ia. 331	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Fluorophenyl
45	Ia. 332	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Fluorophenyl
	Ia. 333	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Fluorophenyl
	Ia. 334	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Methylphenyl

No.	R <sup>1</sup>	R <sup>2</sup>
Ia.335	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Methylphenyl
Ia.336	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Methylphenyl
5 Ia.337	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Methoxyphenyl
Ia.338	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Methoxyphenyl
Ia.339	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Methoxyphenyl
Ia.340	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-(Methoxycarbonyl)phenyl
10 Ia.341	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Methoxycarbonyl)phenyl
Ia.342	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Methoxycarbonyl)phenyl
Ia.343	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Nitrophenyl
Ia.344	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Nitrophenyl
Ia.345	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Nitrophenyl
15 Ia.346	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-(Dimethylamino)phenyl
Ia.347	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Dimethylamino)phenyl
Ia.348	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Dimethylamino)phenyl
Ia.349	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-(Trifluoromethyl)phenyl
20 Ia.350	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Trifluoromethyl)phenyl
Ia.351	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Trifluoromethyl)phenyl
Ia.352	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-(Phenoxy)phenyl
Ia.353	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-(Phenoxy)phenyl
25 Ia.354	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2,4-Difluorophenyl
Ia.355	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2,4-Dichlorophenyl
Ia.356	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,4-Difluorophenyl
Ia.357	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,4-Dichlorophenyl
Ia.358	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,5-Difluorophenyl
30 Ia.359	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3,5-Dichlorophenyl
Ia.360	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Pyridyl
Ia.361	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Pyridyl
Ia.362	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Pyridyl
35 Ia.363	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	α-Naphthyl
Ia.364	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	Benzyl
Ia.365	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Chlorobenzyl
Ia.366	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Chlorobenzyl
40 Ia.367	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Chlorobenzyl
Ia.368	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	2-Methoxybenzyl
Ia.369	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	3-Methoxybenzyl
Ia.370	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	4-Methoxybenzyl
45 Ia.371	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> -Cl
Ia.372	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> -CN
Ia.373	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -CO-OCH <sub>3</sub>


	No.	R <sup>1</sup>	R <sup>2</sup>
	Ia.374	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>
	Ia.375	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> )-CO-OCH <sub>3</sub>
5	Ia.376	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> -OCH <sub>3</sub>
	Ia.377	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.378	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.379	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
10	Ia.380	CH(CH <sub>3</sub> ) <sub>2</sub>	C(CH <sub>3</sub> ) <sub>3</sub>
	Ia.381	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> )-CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>
	Ia.382	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>
	Ia.383	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>
	Ia.384	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -CH=CH <sub>2</sub>
15	Ia.385	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> )=CH <sub>2</sub>
	Ia.386	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> =CH-CH <sub>3</sub>
	Ia.387	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -C≡CH
	Ia.388	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> )-C≡CH
20	Ia.389	CH(CH <sub>3</sub> ) <sub>2</sub>	Cyclopropyl
	Ia.390	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -Cyclopropyl
	Ia.391	CH(CH <sub>3</sub> ) <sub>2</sub>	Cyclopentyl
	Ia.392	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -Cyclopentyl
25	Ia.393	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -(1,3-Dioxolanyl)
	Ia.394	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -(2-Furyl)
	Ia.395	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -(3-Furyl)
	Ia.396	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -(2-Thienyl)
	Ia.397	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>2</sub> -(3-Thienyl)
30	Ia.398	CH(CH <sub>3</sub> ) <sub>2</sub>	Phenyl
	Ia.399	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Chlorophenyl
	Ia.400	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Chlorophenyl
	Ia.401	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Chlorophenyl
35	Ia.402	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Fluorophenyl
	Ia.403	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Fluorophenyl
	Ia.404	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Fluorophenyl
	Ia.405	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Methylphenyl
40	Ia.406	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Methylphenyl
	Ia.407	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Methylphenyl
	Ia.408	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Methoxyphenyl
	Ia.409	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Methoxyphenyl
45	Ia.410	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Methoxyphenyl
	Ia.411	CH(CH <sub>3</sub> ) <sub>2</sub>	2-(Methoxycarbonyl)phenyl
	Ia.412	CH(CH <sub>3</sub> ) <sub>2</sub>	3-(Methoxycarbonyl)phenyl

No.	R <sup>1</sup>	R <sup>2</sup>
Ia.413	CH(CH <sub>3</sub> ) <sub>2</sub>	4-(Methoxycarbonyl)phenyl
Ia.414	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Nitrophenyl
5 Ia.415	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Nitrophenyl
Ia.416	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Nitrophenyl
Ia.417	CH(CH <sub>3</sub> ) <sub>2</sub>	2-(Dimethylamino)phenyl
Ia.418	CH(CH <sub>3</sub> ) <sub>2</sub>	3-(Dimethylamino)phenyl
10 Ia.419	CH(CH <sub>3</sub> ) <sub>2</sub>	4-(Dimethylamino)phenyl
Ia.420	CH(CH <sub>3</sub> ) <sub>2</sub>	2-(Trifluoromethyl)phenyl
Ia.421	CH(CH <sub>3</sub> ) <sub>2</sub>	3-(Trifluoromethyl)phenyl
Ia.422	CH(CH <sub>3</sub> ) <sub>2</sub>	4-(Trifluoromethyl)phenyl
Ia.423	CH(CH <sub>3</sub> ) <sub>2</sub>	3-(Phenoxy)phenyl
15 Ia.424	CH(CH <sub>3</sub> ) <sub>2</sub>	4-(Phenoxy)phenyl
Ia.425	CH(CH <sub>3</sub> ) <sub>2</sub>	2,4-Difluorophenyl
Ia.426	CH(CH <sub>3</sub> ) <sub>2</sub>	2,4-Dichlorophenyl
Ia.427	CH(CH <sub>3</sub> ) <sub>2</sub>	3,4-Difluorophenyl
20 Ia.428	CH(CH <sub>3</sub> ) <sub>2</sub>	3,4-Dichlorophenyl
Ia.429	CH(CH <sub>3</sub> ) <sub>2</sub>	3,5-Difluorophenyl
Ia.430	CH(CH <sub>3</sub> ) <sub>2</sub>	3,5-Dichlorophenyl
Ia.431	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Pyridyl
25 Ia.432	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Pyridyl
Ia.433	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Pyridyl
Ia.434	CH(CH <sub>3</sub> ) <sub>2</sub>	α-Naphthyl
Ia.435	CH(CH <sub>3</sub> ) <sub>2</sub>	Benzyl
Ia.436	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Chlorobenzyl
30 Ia.437	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Chlorobenzyl
Ia.438	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Chlorobenzyl
Ia.439	CH(CH <sub>3</sub> ) <sub>2</sub>	2-Methoxybenzyl
Ia.440	CH(CH <sub>3</sub> ) <sub>2</sub>	3-Methoxybenzyl
35 Ia.441	CH(CH <sub>3</sub> ) <sub>2</sub>	4-Methoxybenzyl
Ia.442	-(CH <sub>2</sub> ) <sub>4</sub> -	
Ia.443	-CH <sub>2</sub> -CH=CH-CH <sub>2</sub> -	

40 Other very especially preferred compounds I are those of the formulae Ib to Iz, Iφ, Iλ, Iπ, Iψ and Iζ, in particular

- the compounds Ib.1 to Ib.443, which differ from the corresponding compounds Ia.1 to Ia.443 only in that R<sup>29</sup> is amino:
- 45


5



Ib;

- 10 - the compounds Ic.1 to Ic.443, which differ from the corresponding compounds Ia.1 to Ia.443 only in that X<sup>1</sup> is hydrogen:


15



Ic;

- 20 - the compounds Id.1 to Id.443, which differ from the corresponding compounds Ia.1 to Ia.443 only in that X<sup>1</sup> is hydrogen and R<sup>29</sup> = amino:

25




Id;

30

- the compounds Ie.1 to Ie.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>5</sup>, A<sup>1</sup> is oxygen, R<sup>7</sup> is difluoromethyl and R<sup>8</sup> is methyl:

35




Ie;

40


- the compounds If.1 to If.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that X<sup>1</sup> is chlorine, Q is Q<sup>5</sup>, A<sup>1</sup> is oxygen, R<sup>7</sup> is difluoromethyl and R<sup>8</sup> is methyl:

45


26




- 10 - the compounds Ig.1 to Ig.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>5</sup>, A<sup>1</sup> is oxygen and R<sup>7</sup> + R<sup>8</sup> is tetramethylene:



- 20 - the compounds Ih.1 to Ih.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that X<sup>1</sup> is chlorine, Q is Q<sup>5</sup>, A<sup>1</sup> is oxygen and R<sup>7</sup> + R<sup>8</sup> is tetramethylene:




- 30 - the compounds Ij.1 to Ij.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>22</sup>, A<sup>10</sup> and A<sup>11</sup> are oxygen, A<sup>12</sup> is sulfur and R<sup>32</sup>, R<sup>33</sup> are methyl:



- 40 - the compounds Ik.1 to Ik.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>22</sup>, A<sup>10</sup>, A<sup>11</sup> & A<sup>12</sup> are oxygen and R<sup>32</sup> & R<sup>33</sup> are methyl:

45


5



Ik,

- 10 - the compounds Im.1 to Im.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>27</sup>, A<sup>13</sup> is oxygen, R<sup>34</sup> & R<sup>36</sup> are hydrogen and R<sup>35</sup> is trifluoromethyl:


15



Im;

- 20 - the compounds In.1 to In.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>27</sup>, A<sup>13</sup> is oxygen, R<sup>34</sup> is hydrogen, R<sup>35</sup> is trifluoromethyl and R<sup>36</sup> is methyl:


25



In;

- 30 - the compounds Io.1 to Io.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>27</sup>, A<sup>13</sup> is oxygen, R<sup>34</sup> is hydrogen, R<sup>35</sup> is SO<sub>2</sub>-CH<sub>3</sub> and R<sup>36</sup> is amino:

35



Io;


- 40 - the compounds Ip.1 to Ip.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>32</sup>, R<sup>37</sup> is chlorine, R<sup>38</sup> is difluoromethoxy and R<sup>39</sup> is methyl:

45



28


5



Ip;

- the compounds Iq.1 to Iq.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>32</sup>, R<sup>37</sup> is bromine, R<sup>38</sup> is difluoromethoxy and R<sup>39</sup> is methyl:


15



Iq;

- the compounds Ir.1 to Ir.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that X<sup>1</sup> is chlorine, Q is Q<sup>32</sup>, R<sup>37</sup> is bromine, R<sup>38</sup> is difluoromethoxy and R<sup>39</sup> is methyl:


25



Ir;

- the compounds Is.1 to Is.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>32</sup>, R<sup>37</sup> is chlorine, R<sup>38</sup> is trifluoromethyl and R<sup>39</sup> is methyl:

35




Is;

- the compounds It.1 to It.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>32</sup>, R<sup>37</sup> is bromine, R<sup>38</sup> is trifluoromethyl and R<sup>39</sup> is methyl:

45

29


5



It;

- the compounds Iu.1 to Iu.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that X<sup>1</sup> is chlorine, Q is Q<sup>32</sup>, R<sup>37</sup> is bromine, R<sup>38</sup> is trifluoromethyl and R<sup>39</sup> is methyl:


15



Iu;

- the compounds Iv.1 to Iv.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>32</sup>, R<sup>37</sup> is chlorine, R<sup>38</sup> is SO<sub>2</sub>-CH<sub>3</sub> and R<sup>39</sup> is methyl:


25



Iv;

- the compounds Iw.1 to Iw.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>32</sup>, R<sup>37</sup> is bromine, R<sup>38</sup> is SO<sub>2</sub>-CH<sub>3</sub> and R<sup>39</sup> is methyl:

35




Iw;


- the compounds Ix.1 to Ix.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that X<sup>1</sup> is chlorine, Q is Q<sup>32</sup>, R<sup>37</sup> is bromine, R<sup>38</sup> is SO<sub>2</sub>-CH<sub>3</sub> and R<sup>39</sup> is methyl:

45


30




- 10 - the compounds Iy.1 to Iy.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>38</sup>, R<sup>40</sup> is chlorine, R<sup>41</sup>, R<sup>43</sup> are hydrogen and R<sup>42</sup> is trifluoromethyl:




- 20 - the compounds Iz.1 to Iz.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>39</sup>, A<sup>1</sup> is oxygen, A<sup>15</sup> is sulfur, R<sup>44</sup> and R<sup>45</sup> are methyl:




- 30 - the compounds Iφ.1 to Iφ.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>40</sup>, A<sup>16</sup> & A<sup>17</sup> are oxygen and R<sup>46</sup> + R<sup>47</sup> form a chain -CH<sub>2</sub>CH<sub>2</sub>-O-CH<sub>2</sub>-:




- 40 - the compounds Iλ.1 to Iλ.443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>40</sup>, A<sup>16</sup> is sulfur, A<sup>17</sup> is oxygen and R<sup>46</sup> + R<sup>47</sup> form a chain -CH<sub>2</sub>CH<sub>2</sub>-O-CH<sub>2</sub>-:




- the compounds I $\pi$ .1 to I $\pi$ .443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>40</sup>, A<sup>16</sup> & A<sup>17</sup> are sulfur and R<sup>46</sup> + R<sup>47</sup> form a chain -CH<sub>2</sub>CH<sub>2</sub>-O-CH<sub>2</sub>-:



- 10
- the compounds I $\psi$ .1 to I $\psi$ .443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>40</sup>, A<sup>16</sup> is oxygen, A<sup>17</sup> is sulfur and R<sup>46</sup> + R<sup>47</sup> form a chain -CH<sub>2</sub>CH<sub>2</sub>-O-CH<sub>2</sub>-:




- 20
- the compounds I $\zeta$ .1 to I $\zeta$ .443, which differ from the corresponding compounds Ia.1 to Ia.443 in that Q is Q<sup>7</sup>, A<sup>3</sup> & A<sup>4</sup> are oxygen and R<sup>9</sup> + R<sup>10</sup> form a tetramethylene chain:



- 30
- The uracil substituted phenyl sulfamoyl carboxamides I according to the invention are obtainable by various routes available and known to those skilled to the art, preferably by one of the processes described hereinbelow.

- 35
- A) Reaction of a benzoic acid derivative II with a sulfamide III, optionally in the presence of a coupling agent such as N,N-carbonyldiimidazole (CDI) or after converting II into the corresponding acid chloride:



N,N'-carbonyldiimidazole (CDI) is added to a solution of the carboxylic acid derivative of formula II in an inert solvent such as tetrahydrofuran. The resulting mixture is stirred under reflux for a sufficient period of time to allow the reaction to come to completion, and is then cooled to room temperature. An optionally substituted sulfamide III is added followed by diazabicycloundecane (DBU) and the mixture is stirred until the reaction is complete. Standard workup and isolation methods give the product in purified form.

The benzoic acid derivatives II - and the corresponding carboxylates, which can be saponified in a simply manner to give the free acids II - are known from the literature or can be prepared analogously to methods known from the literature.

The methods for saponifying the esters to the benzoic acid derivatives II are sufficiently well known to the skilled artisan; consequently, details are not necessary. By way of example, reference is made to Kocienski, "Protecting Groups", Thieme Verlag 1994, and Greene, Wuts, Protecting groups in organic synthesis, Wiley 1999, and Houben-Weyl, Methoden der organischen Chemie, Vol. E5, Part I (1985), pp. 223 et seq.

In addition to activation to the imidazolones other methods are also suitable.

Various methods are suitable for activating the acids. They can, for example, be converted to the acid chloride by treating them with  $\text{SOCl}_2$ ,  $\text{POCl}_3$ ,  $\text{PCl}_5$ ,  $\text{COCl}_2$  or  $(\text{COCl})_2$ . Alternatively, the imidazolidine can be prepared by reaction with N,N'-carbonyldiimidazole. The methods used are sufficiently well known to the skilled artisan, e.g., from Houben Weyl, Methoden der Organischen Chemie, Vol. E5 (1985), Part 1, pp. 587 et seq. and Vol. E5 (1985), Part II, pp. 934 et seq.

Methods of preparing benzoic acid derivatives II where Q is other than  $\text{Q}^{21}$  include those methods described in US 5,872,253, US 5,484,763 and in co-pending patent application Serial Number 09/368,340 filed August 4, 1999 and incorporated herein by reference thereto.

The precursors required for the synthesis of compounds I in which  $\text{Q} = \text{Q}^{21}$ , such as 2-chloro-5[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-4-fluorobenzoic acid (CAS No. 120890-57-5), are described for example in EP-A

195 346, WO 89/02891, WO 98/08151 and the literature cited therein, or may be produced in the manner disclosed therein.


- 5 With regard to the esters of II where  $Q = Q^5$ ,  $A^1 = \text{oxygen}$ ,  $R^7 = \text{difluoromethyl}$ ,  $R^8 = \text{methyl}$ ,  $X^1 = \text{fluorine or chlorine}$  and  $X^2 = \text{chlorine}$ , reference is made to US 5,035,740 and GB-A 22 53 625; with regard to II where  $Q = Q^5$  and where  $R^7$  and  $R^8$  together with the atoms to which they are attached form a 6-membered ring, such as
- 10 2-chloro-4-fluoro-5-(5,6,7,8-tetrahydro-3-oxo-1,2,4-triazolo[4,3-a]pyridin-2(3H)-yl)benzoic acid methyl ester (CAS No. 104799-37-3), reference is made to JP-A 61/069,776. Such compounds are also mentioned in WO 94/22860.
- 15 Benzoic acid derivatives II where  $Q$  is  $Q^{22}$ ,  $A^{10} \& A^{11} = \text{oxygen}$ ,  $A^{12} = \text{oxygen or sulfur}$ ,  $R^{32} \& R^{33} = \text{amino or alkyl}$ ,  $X^1 = \text{fluorine}$  and  $X^2 = \text{chlorine}$  are known from EP-A 584,655 and WO 00/50409, e.g. 2-chloro-5-(3,5-dimethyl-2,6-dioxo-4-thi-oxo-1,3,5-triazinan-1-yl)-4-fluorobenzoic acid (CAS No. 289882-59-3) and 2-chloro-5-(3,5-dimethyl-2,4,6-trioxo-1,3,5-triazinan-1-yl)-4-fluorobenzoic acid methyl ester (CAS No. 154883-47-3).
- 20 Benzoic acid derivatives II and their esters where  $Q$  is  $Q^{27}$  are known from WO 97/07104, WO 96/39392, WO 99/14201 and WO 99/52878, e.g. 2-chloro-4-fluoro-5-(5-trifluormethyl-3-pyridazinon-2-yl) benzoic acid ( $R^{34} = \text{hydrogen}$ ,  $R^{35} = \text{trifluoromethyl}$ ,  $R^{36} = \text{hydrogen}$ ,  $X^1 = \text{fluorine}$  and  $X^2 = \text{chlorine}$ ) and 2-chloro-4-fluoro-5-(4-trifluoromethyl-5-trifluoromethyl-3-pyridazinon-2-yl) benzoic acid (CAS No. 259141-58-7;  $R^{34} = \text{hydrogen}$ ,  $R^{35} = \text{trifluoromethyl}$ ,  $R^{36} = \text{methyl}$ ,  $X^1 = \text{fluorine}$ ,  $X^2 = \text{chlorine}$ ).
- 30 Benzoic acid derivatives II where  $Q$  is  $Q^{32}$  are known from EP-A 361,114, WO 92/06962, WO 96/02515, US 6,096,689 and WO 98/38169, e.g. 4-Chloro-3-[4-chloro-2-fluoro-5-carboxyphenyl]-5-difluoromethoxy-1-methyl-1H-pyrazole (CAS No. 129631-53-4;  $Q = Q^{32}$ ,  $R^{37} = \text{chlorine}$ ,  $R^{38} = \text{difluoromethoxy}$ ,  $R^{39} = \text{methyl}$ ,  $X^1 = \text{fluorine}$ ,  $X^2 = \text{chlorine}$ ), 4-Chloro-3-[4-chloro-2-fluoro-5-carboxyphenyl]-5-trifluoromethyl-1-methyl-1H-pyrazole (CAS-No. 142622-56-8;  $Q = Q^{32}$ ,  $R^{37} = \text{chlorine}$ ,  $R^{38} = \text{difluoromethoxy}$ ,  $R^{39} = \text{methyl}$ ,  $X^1 = \text{fluorine}$ ,  $X^2 = \text{chlorine}$ ), or can be prepared in a manner similar to that described there.
- 45

Benzoic acid derivatives II where Q is Q<sup>38</sup> are known from WO 95/02580, US 5,783,522 and WO 98/07700, e.g. 2-chloro-5-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-4-fluorobenzoic acid (CAS No. 188782-31-2), or can be prepared in a manner similar to that described there.

Benzoic acid derivatives II where Q is Q<sup>39</sup> are known from WO 99/59983 and DE-A 19 835 943, or can be prepared in a manner similar to that described there.

Benzoic acid derivatives II where Q is Q<sup>40</sup> are known from WO 94/10173 and WO 00/01700, or can be prepared in a manner similar to that described there.

Benzoic acid derivatives II where Q is Q<sup>5</sup> can be prepared according to US 5,035,740 as follows:




The hydrazines IV are known, e.g, from WO 97/07104 ( $\text{X}^1 = \text{fluorine}$ ), or may be prepared in known manner.

The sulfamides of the formula III are obtainable according to methods known per se, for example analogously to the method described in Hamprecht et al., Angew. Chemie 93, 151 (1981) and Houben-Weyl, Methoden der Organischen Chemie, Vol. E11 (1985), pp. 1019 et seq.

As an example, formula III sulfamides where  $\text{X}^3$  is hydrogen may be prepared by reaction of S-chlorosulfonamide with an amine  $\text{HNR}^1\text{R}^2$ :




36



Formula III sulfamides where  $\text{X}^3$  is not hydrogen may be prepared by reaction of sulfonyl chloride with an amine  $\text{HNR}^1\text{R}^2$  to give the sulfamoyl chloride compound  $\text{Cl}-\text{SO}_2-\text{NR}^1\text{R}^2$ , and reacting said sulfamoyl chloride compound with an amine  $\text{X}^3-\text{NH}_2$ :



B) Displacement of a halide by Q:




Hal = halogen, preferably fluorine, chlorine or bromine.

By this route, an aniline of formula IV is converted to a diazonium salt, then treated with iodine and potassium iodide to give the iodo compound of the formula V. Reaction of formula V compounds with an unsubstituted QH moiety, for example, a uracil of formula  $\text{Q}^{21}\text{H}$  in the presence of a copper(I) catalyst gives a final product of formula Ia. In this way, compounds I according to the invention where  $\text{Q} = \text{Q}^{21}$  can be obtained, by analogy to the method disclosed by T. Maruyama, K. Fujiwara and M. Fukuhara in J. Chem. Soc., Perkin Trans. 1995 (7), pp. 733-734, where Hal = iodine, and the reaction is carried out with the addition of a Cu(I) source.

However, transition-metal-free methods are also suitable if the substituents Hal,  $\text{X}^1$  and  $\text{X}^2$  are properly selected. In this respect, reference is made by way of example to WO 96/39392, which describes methods which are suitable for the manufacture of compounds I where  $\text{Q} = \text{Q}^{27}$ .

The haloaryl precursors V can be obtained by a Sandmeyer reaction from the corresponding anilines (see also formula scheme V). These methods are sufficiently well known to the skilled artisan, so reference is only made here to Houben-Weyl, Methoden der Org. Chemie, Vol. 5/4, 4th edition 1960, pp. 438 et seq.


- C) Reaction of an aniline intermediate VI with an oxazinone compound of the formula VII to give a compound I where A is oxygen, X<sup>3</sup> is hydrogen, Q is Q<sup>21</sup>, A<sup>8</sup> & A<sup>9</sup> are oxygen and R<sup>29</sup> is hydrogen, optionally followed by alkylation and hydrolysis:



R<sup>29</sup>-Hal represents a C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-haloalkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>2</sub>-C<sub>6</sub>-alkenyl, C<sub>2</sub>-C<sub>6</sub>-haloalkenyl or C<sub>3</sub>-C<sub>6</sub>-alkynyl halide.

Among the methods known for the preparation of oxazinone compounds VII are those described in WO 99/14216.

Formula VI aniline derivatives may be prepared by conventional procedures such as the conversion of the appropriately substituted benzoic acid IX to the corresponding sulfamoyl carboxamide X (see method A, above), which in turn is then nitrated and reduced:



Suitable nitration reagents are for example nitric acid in various concentrations, including concentrated and fuming nitric acid, mixtures of sulfuric and nitric acid, and acetyl nitrates and alkyl nitrates.

The reaction can be carried out either without a solvent in an excess of the nitration reagent or in an inert solvent or diluent, suitable agents being, for example, water, mineral acids, organic acids, halohydrocarbons such as methylene chloride, anhydrides such as acetic anhydride, and mixtures thereof.

The sulfamoyl carboxamide X and the nitration reagent are expediently employed in approximately equimolar amounts; with regard to the yield of X, it may be advantageous to use the nitration reagent in an excess of up to about 10 times the molar amount, based on the amount of X. When the reaction is carried out without a solvent in the nitration reagent, the latter is present in an even greater excess.

The reaction temperature is generally from  $(-100)^{\circ}\text{C}$  to  $200^{\circ}\text{C}$ , preferably from  $(-30)$  to  $50^{\circ}\text{C}$ .

The nitrated compounds XI can then be reduced to the aniline derivatives VI.

The reduction is generally carried out by reaction of the nitro compound with a transition metal such as iron, zinc or tin under acidic conditions or with a complex hydride such as

lithium aluminium hydride and sodium borohydride, the reduction being carried out in bulk or in a solvent or diluent.

- 5 Examples of suitable solvents are - depending on the reducing agent selected - water, alcohols such as methanol, ethanol and isopropanol, or ethers such as diethyl ether, methyl tert-butyl ether, dioxane, tetrahydrofuran and ethylene glycol dimethyl ether.
- 10 If a metal is used for reduction purposes, it is preferable to work without a solvent in an inorganic acid, especially in concentrated or dilute hydrochloric acid, or in a liquid organic acid such as acetic acid and propionic acid. However,
- 15 the acid can also be diluted with an inert solvent, e.g., one of those mentioned above. The reduction with complex hydrides is carried out preferably in a solvent, for example an ether or an alcohol.
- 20 The nitrated compound XI and the reducing agent are frequently used in approximately equimolar amounts; to optimize the reaction it may however be advantageous to use either component in an excess of up to about the 10-fold molar amount.
- 25 The amount of acid is not critical. So as to reduce the starting compound as completely as possible, it is expedient to use at least an equivalent amount of acid. Frequently, the acid is used in excess, based on the nitrated compound XI.
- 30 The reaction temperature is generally from (-30) to 200°C, preferably from 0 to 80°C.
- 35 For working up, the reaction mixture is as a rule diluted with water and the product is isolated by filtration, crystallization or extraction with a solvent which is substantially immiscible with water, e.g., ethyl acetate, diethyl ether or methylene chloride. If desired, the product VI can then be purified in conventional manner.
- 40 The nitro group of compounds XI can also be hydrogenated catalytically with hydrogen. Examples of suitable catalysts to this end are Raney nickel, palladium on charcoal, palladium oxide, platinum and platinum oxide. An amount of from 0.05 to 50 mol%, based on the compound XI to be reduced,
- 45 is generally sufficient.

It is possible to dispense with a solvent, or to use an inert solvent or diluent, e.g., acetic acid, a mixture of acetic acid and water, ethyl acetate, ethanol or toluene. When the catalyst has been separated off, the reaction solution can be worked up as usual to give the product VI. Hydrogenation can be effected at atmospheric or superatmospheric hydrogen pressure.

Further methods and reaction conditions are given in the literature (see, for example, Houben-Weyl, Methoden der Organischen Chemie, nitrogen compounds I, Part 1 (1971), Vol. X/1, pp. 463 et seq.).

Not only the compounds I according to the invention where  $Q = Q^{21}$ , but also compounds I where  $Q = Q^7$ ,  $Q^{22}$  or  $Q^{40}$  can be produced from the aniline derivatives VI. To prepare compounds I where  $Q = Q^{22}$ , reference is made to the methods described in WO 00/50409 and EP-A 584 655, and to prepare compounds I where  $Q = Q^{40}$ , reference is made to the methods taught in WO 94/10173 and WO 00/01700.


The aniline derivatives VI can, however, also be converted in conventional manner (see, for example, WO 97/07104 and Houben-Weyl, Methoden der Organischen Chemie, Vol. E1, nitrogen compounds) to the corresponding hydrazines, from which compounds I where  $Q = Q^5$  oder  $Q^{27}$  can be prepared.

Further methods for preparing compounds I according to the invention are given in Böger, Wakabayashi Peroxidizing herbicides, Springer Verlag 1999.

D) Reacting a benzoic acid derivative VIII with an electrophilic amination reagent in the presence of a base to give the corresponding N-amino uracil benzoic ester, hydrolyzing said ester to give the benzoic acid II (with  $Q = Q^{21}$ ;  $A^8$  &  $A^9 = O$ ;  $R^{29} = NH_2$ ) and converting the latter to the compounds I ( $A = O$ ;  $Q = Q^{21}$ ;  $A^8$  &  $A^9 = O$ ;  $R^{29} = NH_2$ ) by the route described above:

40

45



25 Examples of electrophilic amination reagents are in particular 2,4-dinitrophenylhydroxylamine and O-mesitylene-sulfonyl hydroxylamine.

Examples of suitable reaction conditions are given in DE-A 19 652 431.

30 All the processes described above are expediently carried out under atmospheric pressure or under the inherent pressure of the reaction mixture in question.

35 As a rule, the reaction mixtures are worked up by methods known per se, for example by removing the solvent, partitioning the residue between a mixture of water and a suitable organic solvent and working up the organic phase to obtain the product.

40 The uracil substituted phenyl sulfamoyl carboxamides I according to the invention can be obtained from the preparation as isomer mixtures which, if desired, can be separated into the pure isomers by the methods conventionally used for this purpose, eg. by means of crystallization or chromatography on an optically active adsorbate. Pure optically active isomers can, for example, 45 also be prepared from suitable optically active starting materials.

Compounds I with C-H acidic substituents can be converted into their alkali metal salts in a manner known per se by reaction with a base of the corresponding cation.

- 5 Salts of I whose metal ion is not an alkali metal ion can normally be prepared by double decomposition of the corresponding alkali metal salt in aqueous solution.

Other metal salts, such as manganese, copper, zinc, iron,  
10 calcium, magnesium and barium salts, can be prepared from the sodium salts in the customary manner, and also ammonium and phosphonium salts by means of ammonia, phosphonium, sulfonium or sulfoxonium hydroxides.

- 15 The compounds I and their agriculturally useful salts are suitable as herbicides, both in the form of isomer mixtures and in the form of the pure isomers. The herbicidal compositions comprising I effect very good control of vegetation on non-crop areas, especially at high rates of application. In crops such as  
20 wheat, rice, maize, soybeans and cotton they act against broad-leaved weeds and grass weeds without damaging the crop plants substantially. This effect is observed especially at low rates of application.

- 25 Depending on the application method in question, the compounds I, or compositions comprising them, can additionally be employed in a further number of crop plants for eliminating undesirable plants. Examples of suitable crops are the following:

Allium cepa, Ananas comosus, Arachis hypogaea, Asparagus  
30 officinalis, Beta vulgaris spec. altissima, Beta vulgaris spec. rapa, Brassica napus var. napus, Brassica napus var. napobrassica, Brassica rapa var. silvestris, Camellia sinensis, Carthamus tinctorius, Carya illinoensis, Citrus limon, Citrus sinensis, Coffea arabica (Coffea canephora, Coffea liberica),  
35 Cucumis sativus, Cynodon dactylon, Daucus carota, Elaeis guineensis, Fragaria vesca, Glycine max, Gossypium hirsutum, (Gossypium arboreum, Gossypium herbaceum, Gossypium vitifolium), Helianthus annuus, Hevea brasiliensis, Hordeum vulgare, Humulus lupulus, Ipomoea batatas, Juglans regia, Lens culinaris, Linum  
40 usitatissimum, Lycopersicon lycopersicum, Malus spec., Manihot esculenta, Medicago sativa, Musa spec., Nicotiana tabacum (N. rustica), Olea europaea, Oryza sativa, Phaseolus lunatus, Phaseolus vulgaris, Picea abies, Pinus spec., Pisum sativum, Prunus avium, Prunus persica, Pyrus communis, Ribes sylvestre,  
45 Ricinus communis, Saccharum officinarum, Secale cereale, Solanum tuberosum, Sorghum bicolor (s. vulgare), Theobroma cacao, Trifolium pratense, Triticum aestivum, Triticum durum, Vicia

faba, Vitis vinifera and Zea mays.

Moreover, the compounds I may also be used in crops which have been made fully or partially tolerant to the action of herbicides 5 due to breeding including genetic engineering methods.

Furthermore, the substituted hydroximic acid derivatives I are also suitable for the desiccation and/or defoliation of plants.

10 As desiccants, they are especially suitable for desiccating the aerial parts of crop plants such as potatoes, oilseed rape, sunflowers and soybeans. This allows completely mechanical harvesting of these important crop plants.

15 Also of economic interest is facilitated harvesting, which is made possible by concentrating, over a period of time, dehiscence, or reduced adhesion to the tree, in the case of citrus fruit, olives or other species and varieties of pomaceous fruit, stone fruit and nuts. The same mechanism, ie. promotion of 20 the formation of abscission tissue between fruit or leaf and shoot of the plants, is also essential for readily controllable defoliation of useful plants, in particular cotton.

Moreover, a shortened period of time within which the individual 25 cotton plants ripen results in an increased fiber quality after harvesting.

The compounds I, or the compositions comprising them, can be employed, for example, in the form of directly sprayable aqueous. 30 solutions, powders, suspensions, also highly-concentrated aqueous, oily or other suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading or granules, by means of spraying, atomizing, dusting, spreading or pouring. The use forms depend on the intended purposes; in any case, they 35 should guarantee the finest possible distribution of the active ingredients according to the invention.

Suitable inert additives are essentially: mineral oil fractions of medium to high boiling point such as kerosene and diesel oil, 40 furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, eg. paraffins, tetrahydronaphthalene, alkylated naphthalenes and their derivatives, alkylated benzenes and their derivatives, alcohols such as methanol, ethanol, propanol, butanol and cyclohexanol, 45 ketones such as cyclohexanone or strongly polar solvents, eg. amines such as N-methylpyrrolidone or water.



Aqueous use forms can be prepared from emulsion concentrates, suspensions, pastes, wettable powders or water-dispersible granules by adding water. To prepare emulsions, pastes or oil dispersions, the substituted hydroximic acid derivatives as such  
5 or dissolved in an oil or solvent, can be homogenized in water by means of wetting agent, tackifier, dispersant or emulsifier. However, it is also possible to prepare concentrates composed of active substance, wetting agent, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and these  
10 concentrates are suitable for dilution with water.

Suitable surfactants are the alkali metal, alkaline earth metal and ammonium salts of aromatic sulfonic acids, eg. ligno-, phenol-, naphthalene- and dibutyl-naphthalenesulfonic acid, and of  
15 fatty acids, of alkyl- and alkylaryl sulfonates, of alkyl sulfates, lauryl ether sulfates and fatty alcohol sulfates, and salts of sulfated hexa-, hepta- and octadecanols, and of fatty alcohol glycol ether, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene, or  
20 of the naphthalenesulfonic acids, with phenol and formaldehyde, polyoxyethylene octylphenyl ether, ethoxylated isooctyl-, octyl- or nonylphenol, alkylphenyl and tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil,  
25 polyoxyethylene alkyl ethers or polyoxypropylene alkyl ethers, lauryl alcohol polyglycol ether acetate, sorbitol esters, lignin-sulfite waste liquors or methylcellulose.

Powders, materials for spreading and dusts can be prepared by  
30 mixing or concomitantly grinding the active substances with a solid carrier.

Granules, eg. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active  
35 ingredients to solid carriers. Solid carriers are mineral earths such as silicas, silica gels, silicates, talc, kaolin, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate,  
40 ammonium phosphate, ammonium nitrate, ureas and products of vegetable origin such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders or other solid carriers.

The concentrations of the active ingredients I in the  
45 ready-to-use products can be varied within wide ranges. In general, the formulations comprise approximately from 0.001 to 98% by weight, preferably 0.01 to 95% by weight, of at least one

active ingredient. The active ingredients are normally employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

5 The formulation examples below illustrate the preparation of such formulations:

- 10 I. 20 parts by weight of an uracil substituted phenyl sulfamoyl carboxamide I are dissolved in a mixture composed of 80 parts by weight of alkylated benzene, 10 parts by weight of the adduct of 8 to 10 mol of ethylene oxide and 1 mol of oleic acid N-monoethanolamide, 5 parts by weight of calcium dodecylbenzenesulfonate and 5 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil. 15 Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.
- 20 II. 20 parts by weight of an uracil substituted phenyl sulfamoyl carboxamide I are dissolved in a mixture composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient. 25 30
- III. 20 parts by weight of an uracil substituted phenyl sulfamoyl carboxamide I are dissolved in a mixture composed of 25 parts by weight of cyclohexanone, 65 parts by weight of a mineral oil fraction of boiling point 210 to 280°C and 10 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient. 35 40
- IV. 20 parts by weight of an uracil substituted phenyl sulfamoyl carboxamide I are mixed thoroughly with 3 parts by weight of sodium diisobutyl naphthalene- $\alpha$ -sulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel and the mixture is ground in a 45

hammer mill. Finely distributing the mixture in 20,000 parts by weight of water gives a spray mixture which comprises 0.1% by weight of the active ingredient.

- 5 V. 3 parts by weight of an uracil substituted phenyl sulfamoyl carboxamide I are mixed with 97 parts by weight of finely divided kaolin. This gives a dust which comprises 3% by weight of the active ingredient.
- 10 VI. 20 parts by weight of an uracil substituted phenyl sulfamoyl carboxamide I are mixed intimately with 2 parts by weight of calcium dodecylbenzenesulfonate, 8 parts by weight of fatty alcohol polyglycol ether, 2 parts by weight of the sodium salt of a phenol/urea/formaldehyde condensate and 68 parts by weight of a paraffinic mineral oil. This gives a stable oily dispersion.
- 15
- VII. 1 part by weight of an uracil substituted phenyl sulfamoyl carboxamide I is dissolved in a mixture composed of 70 parts by weight of cyclohexanone, 20 parts by weight of ethoxylated isooctylphenol and 10 parts by weight of ethoxylated castor oil. This gives a stable emulsion concentrate.
- 20
- 25 VIII. 1 part by weight of an uracil substituted phenyl sulfamoyl carboxamide I is dissolved in a mixture composed of 80 parts by weight of cyclohexanone and 20 parts by weight of Wettol® EM 31 (non-ionic emulsifier based on ethoxylated castor oil; BASF AG). This gives a stable emulsion concentrate.
- 30

The active ingredients I, or the herbicidal compositions comprising them, can be applied pre- or post-emergence. If the active ingredients are less well tolerated by certain crop plants, application techniques may be used in which the herbicidal compositions are sprayed, with the aid of the spray apparatus, in such a way that they come into as little contact as possible, if any, with the leaves of the sensitive crop plants while reaching the leaves of undesirable plants which grow underneath, or the bare soil (post-directed, lay-by).

Depending on the intended aim of the control measures, the season, the target plants and the growth stage, the application rates of active ingredient are from 0.001 to 3.0, preferably 0.01 to 1 kg/ha active substance (a.s.).

- To widen the spectrum of action and to achieve synergistic effects, the substituted hydroximic acid derivatives I can be mixed and applied jointly with a large number of representatives of other groups of herbicidally or growth-regulatory active ingredients. Suitable components in mixtures are, for example,
- 5 1,2,4-thiadiazoles, 1,3,4-thiadiazoles, amides, aminophosphoric acid and its derivatives, aminotriazoles, anilides, (het)aryloxyalkanoic acids and their derivatives, benzoic acid and its derivatives, benzothiadiazinones,
  - 10 2-aryl-1,3-cyclohexanediones, hetaryl aryl ketones, benzylisoxazolidinones, meta-CF<sub>3</sub>-phenyl derivatives, carbamates, quinolinecarboxylic acid and its derivatives, chloroacetanilides, cyclohexane-1,3-dione derivatives, diazines, dichloropropionic acid and its derivatives, dihydrobenzofurans,
  - 15 dihydrofuran-3-ones, dinitroanilines, dinitrophenols, diphenyl ethers, dipyridyls, halocarboxylic acids and their derivatives, ureas, 3-phenyluracils, imidazoles, imidazolinones, N-phenyl-3,4,5,6-tetrahydrophthalimides, oxadiazoles, oxiranes, phenols, aryloxy- or hetaryloxyphenoxypropionic esters,
  - 20 phenylacetic acid and its derivatives, phenylpropionic acid and its derivatives, pyrazoles, phenylpyrazoles, pyridazines, pyridinecarboxylic acid and its derivatives, pyrimidyl ethers, sulfonamides, sulfonylureas, triazines, triazinones, triazolinones, triazolecarboxamides and uracils.

25


- Moreover, it may be advantageous to apply the compounds I, alone or in combination with other herbicides, in the form of a mixture with additional other crop protection agents, for example with pesticides or agents for controlling phytopathogenic fungi or
- 30 bacteria. Also of interest is the miscibility with mineral salt solutions which are employed for treating nutritional and trace element deficiencies. Non-phytotoxic oils and oil concentrates can also be added.

35

#### PREPARATION EXAMPLES


- In order to facilitate a further understanding of the invention, the following examples are presented to illustrate more specific details thereof. The term NMR designates nuclear magnetic resonance; HPLC designates high performance liquid
- 40 chromatography; TLC designates thin layer chromatography; GLC designates gas-liquid chromatography and IR designates infrared spectroscopy.

## Example 1: Preparation of 2-Chloro-4-fluoro-5-nitrobenzoic Acid



A solution of 2-chloro-4-fluorobenzoic acid (24.4 g, 0.142 mol) in 150 ml of concentrated sulfuric acid at 0°C was treated dropwise with 90% nitric acid (13.2 ml, 20 mol%, 0.284 mol) over a 10 min. period at 10°C, stirred for 2.5 hours at 0 to 10°C, poured onto one liter of ice. The white solid was filtered. The filtercake was air-dried and recrystallized from ethyl acetate/heptane to afford the title compound as off-white needles. Yield: 18.0 g (58.1%); identified by NMR spectral analysis.


## 15 EXAMPLE 2: Preparation of 2-Chloro-4-fluoro-5-aminobenzoic acid



20 A solution of 2-chloro-4-fluoro-5-nitrobenzoic acid (18.0 g, 0.0824 mol) in 75 ml of acetic acid was heated at reflux temperature. Iron powder (18.4 g, 0.328 mol) was added in several portions and the resulting suspension was cooled to room temperature and diluted with water and ethyl acetate. The mixture was filtered and the filtrate was saved. The organic layer was washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound as a tan solid. Yield: 9.00 g (58.1%); mp.: 153-155°C; identified by NMR and mass spectral analysis.

30

## EXAMPLE 3: Preparation of 3-(5-Carboxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-6-trifluoromethylpyrimidin-2,4-dione




40 A mixture of 2-chloro-4-fluoro-5-aminobenzoic acid (8.30 g, 0.04308 mol), 2-dimethylamino-4-(trifluoromethyl)-6H-1,3-oxazin-6-one (9.57 g, 0.0460 mol) and acetic acid was stirred three hours at reflux temperature, diluted with ice water and extracted twice with ethyl acetate. The combined organic layers were washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound as a tan solid. Yield: 14.0 g (92.1%); identified by NMR and mass

spectral analysis.

Example 4: Preparation of 3-(5-Carboxymethoxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-methyl-6-trifluoromethylpyrimidin-2,4-dione

5

10




A mixture of 3-(5-carboxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-6-trifluoromethylpyrimidin-2,4-dione (13.3 g, 0.0377 mol), potassium carbonate (13.0 g, 0.0943 mol), methyl iodide (5.87 ml, 0.0943 mol) and dimethyl formamide (150 ml) was stirred overnight at room temperature and diluted with water (500 ml). The resulting mixture was extracted three times with ethyl acetate. The combined organic layers were washed three times with water, aqueous sodium hydroxide (0.1 N) and brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give a beige solid. Recrystallization of the residue from ethanol-water (250 ml) afforded the title compound as white needles. Yield: 11.5 g (80.4%); mp.: 172-173°C; identified by NMR and mass spectral analysis.

Example 5: Preparation of 3-(5-Carbomethoxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-6-trifluoromethylpyrimidin-2,4-dione

30


35



Carbonyl diimidazole (2.97 g, 18.4 mmol) was added to a solution of 3-(5-carboxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-6-trifluoromethylpyrimidin-2,4-dione (4.61 g, 13.1 mmol) in tetrahydrofuran and the resulting mixture was heated to reflux temperature, stirred two minutes and cooled to room temperature. Methanol (2.70 ml, 66.6 mmol) was added and the mixture was stirred overnight at room temperature. Subsequently, the mixture was concentrated under reduced pressure and the resultant residue was taken up in methylene chloride. The organic mixture was washed twice with hydrochloric acid (10% aqueous and 5% aqueous) and water. The organic layer was concentrated under reduced


pressure to give a brown solid, which was suspended in methylene chloride, followed by filtration. The filtercake was washed three times with methylene chloride, filtered and dried to afford the title compound as a white solid, which was identified by NMR spectral analysis. Yield: 4.27 g (89.0%).

EXAMPLE 6: Preparation of 3-(5-Carbomethoxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-amino-6-trifluoromethylpyrimidin-2,4-dione



To a suspension of 3-(5-carbomethoxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-6-trifluoromethylpyrimidin-2,4-dione (4.24 g, 11.6 mmol) in anhydrous tetrahydrofuran was added potassium carbonate (1.60 g, 11.6 mmol) followed by O-mesitylenesulfonyl hydroxylamine (3.04 g, 14.1 mmol; J.G. Krause, *Synthesis*, 1972, 140). The resulting mixture was stirred overnight at room temperature and diluted with water. The mixture was extracted four times with ethyl acetate. The combined extracts were dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound as a foam, which was identified by NMR spectral analysis. Yield: 4.63 g (>100%).


EXAMPLE 7: Preparation of 3-(5-Carboxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-amino-6-trifluoromethylpyrimidin-2,4-dione



To a solution of 3-(5-carbomethoxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-amino-6-trifluoromethylpyrimidin-2,4-dione (1.53 g, 4.01 mmol) in anhydrous methylene chloride was added boron tribromide (1M in methylene chloride, 16.0 ml, 16.0 mmol).  
5 The resultant mixture was stirred overnight at room temperature and then diluted with water. The aqueous layer was separated and allowed to stand at room temperature overnight; filtration and drying afforded the title compound as a white solid, which was identified by NMR and mass spectral analysis. Yield: 0.61 g  
10 (41.5%).


The original organic layer was concentrated under reduced pressure to a glassy solid, which was triturated with water to afford an additional amount of the title compound as a tan solid, which was identified by NMR and mass spectral analysis. Yield:  
15 0.310 g (21.1%); mp.: 150°C (decomposition).

EXAMPLE 8: Preparation of 3-(5-carboxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-methyl-6-trifluoromethylpyrimidin-2,4-dione



Boron tribromide (84.0 ml, 0.0840 mol, 1M in methylene chloride) was added dropwise to a mixture of 3-(5-carbomethoxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-methyl-6-trifluoromethylpyrimidin-2,4-dione (10.7 g, 0.0281 mol) and methylene chloride (150 ml). The resulting mixture was stirred overnight at room temperature and diluted with ice water. The organic layer is saved and the aqueous layer was extracted twice with ethyl acetate. The extracts were combined with the organic layer,  
35 washed with brine, dried over anhydrous magnesium sulfate and concentrated under reduced pressure to afford the title compound as a white solid, which was identified by NMR and mass spectral analysis. Yield: 10.2 g (100%); mp.: 240-241°C.

40 EXAMPLE 9: Preparation of S-Chlorosulfonamide




45 Formic acid (10.8 ml, 0.287 mol) was added dropwise over a two hour period to chlorosulfonyl isocyanate (25.0 ml, 0.287 mol), maintaining the temperature below 20°C. The resulting suspension



was stirred two hours at 20°C and diluted with anhydrous toluene (100 ml). The resulting mixture was stirred overnight at ambient temperature and filtered. The filtrate was concentrated under reduced pressure to afford the title compound as an off-white solid, which was identified by IR spectral analysis. Yield: 32.1 g (97.3%).

EXAMPLE 10: Preparation of N-Methylsulfamide



A solution of S-chlorosulfonamide (3.00 g, 0.0260 mol) in tetrahydrofuran (10 ml) was added dropwise to methylamine (40 ml, 2M in tetrahydrofuran) at 10°C. The resulting mixture was stirred one hour at 0°C and three days at room temperature. The suspension was filtered and the filtrate concentrated under reduced pressure to give a yellow solid. Chromatography on silica gel (9:1 methylene chloride - methanol) afforded the title compound as an off-white solid, which was identified by NMR and mass spectral analysis. Yield: 1.36 g (47.4%).


EXAMPLES 11-15: Preparation of N-substituted sulfamides

25 Using essentially the same procedure as described in Example 10 hereinabove and substituting the appropriate amine starting material, the following compounds are prepared:




Example	R <sup>1</sup>	R <sup>2</sup>	mp. [°C]
11	H	CH <sub>2</sub> -C(CH <sub>3</sub> ) <sub>3</sub>	—
12	CH <sub>3</sub>	CH <sub>2</sub> -CH=CH <sub>2</sub>	36-38
13	CH <sub>3</sub>	benzyl	91-94
14	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	—
15	H	H	—

40 Example 16: Preparation of O-2,4,5-trichlorophenyl sulfamate



A solution of 2,4,5-trichlorophenol (96.0 g, 0.486 mol) in toluene (95 ml) was treated dropwise with chlorosulfonyl isocyanate (67.4 g, 0.476 mol) at 40-55°C. The resulting mixture was stirred three hours at reflux temperature, cooled to 40°C and  
 5 quenched with water until gas evolution ceases. The suspension was filtered and the filtercake was air-dried to afford the title compound as a white solid, which is identified by NMR and IR spectral analysis. Yield: 118.5 g (90.0%).

10 Example 17: Preparation of N-methyl-N-isopropyl sulfamide



Triethylamine (2.50 ml, 0.0181 mol) was added to a solution of methyl propargylamine (1.55 ml, 0.0181 mol) in acetonitrile. To  
 20 the resulting mixture was added 0-2,4,5-trichlorophenyl sulfamate (5.00 g, 0.0181 mol). The resulting mixture was stirred for one hour at room temperature and filtered through silica gel with methylene chloride. The filtrate was concentrated under reduced pressure to afford a white solid. Chromatography of the residue  
 25 on silica gel (0.5% methanol-methylene chloride) afforded the title compound, which was identified by NMR spectral analysis. Yield: 1.84 g (68.7%).

EXAMPLES 18-52: Preparation of N-substituted sulfamides

30

Using essentially the same procedure as described in Example 17 hereinabove and substituting the appropriate amine starting material, the following sulfamides were prepared:




Example	R <sup>1</sup>	R <sup>2</sup>	mp. [°C] / <sup>1</sup> H-NMR [ppm]
18	CH <sub>3</sub>	CH <sub>3</sub>	—
19	CH <sub>3</sub>	3-chlorobenzyl	—
40 20	CH <sub>3</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	—
21	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	39-41
22	H	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>	—
23	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	32-34
45 24	H	C(CH <sub>3</sub> ) <sub>3</sub>	49-53
25	H	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	32-35
26	CH <sub>3</sub>	3-methoxybenzyl	—

## 54

Example	R <sup>1</sup>	R <sup>2</sup>	mp. [°C] / <sup>1</sup> H-NMR [ppm]
27	CH <sub>3</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	103-104
28	H	CH(CH <sub>3</sub> ) <sub>2</sub>	—
29	CH <sub>3</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	63-65
30	H	C <sub>2</sub> H <sub>5</sub>	—
31	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	—
32	H	CH <sub>2</sub> -CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	—
33	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -phenyl	—
34	CH <sub>3</sub>	phenyl	84-86
35	H	CH <sub>2</sub> -C≡CH	—
36	H	CH <sub>2</sub> -(2-furyl)	62-64
37	CH <sub>3</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>	—
38	H	CH <sub>2</sub> -(2-thienyl)	93-96
39	H	cyclopentyl	55
40	H	4-methoxybenzyl	—
41	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	—
42	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -CN	—
43	CH <sub>3</sub>	CH <sub>2</sub> -(1,3-dioxalanyl)	—
44	H	4-chlorobenzyl	—
45	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>	54-57
46		-CH <sub>2</sub> -CH=CH-CH <sub>2</sub> -	—
47	H	cyclopropyl	58-60
48		-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -	84-86
49	CH <sub>3</sub>	cyclopropyl	—
50	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	—
51	H	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	—
52	H	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	—
53	CH <sub>3</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>	5.1(br., s, 2H), 4.25(q, 2H), 4.1(s, 2H), 3.0(s, 3H), 1.3(t, 3H)

EXAMPLE 54: Preparation of 3-(5-(N,N-dimethyl)sulfamoylcarbox-  
amido-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-6-tri-  
fluoromethylpyrimidin-2,4-dione



To a solution of 3-(5-carboxy-4-chloro-2-fluorophenyl)-1,2,3,4-dihydro-1-methyl-6-(trifluoromethyl)pyrimidin-2,4-dione (1.50 g, 4.09 mmol) in tetrahydrofuran was added N,N'-carbonyldiimidazole (1.00 g, 6.14 mmol). The resulting mixture was stirred one hour at reflux temperature and cooled to room temperature. Dimethyl sulfamide (0.760 g, 6.14 mmol) was added, followed by diazabicycloundecane (0.930 ml, 6.14 mmol) after 10 min. The resulting mixture was stirred overnight at room temperature and then concentrated under reduced pressure. The resultant residue was

20

25

30

partitioned between ethyl acetate and hydrochloric acid (2N). The organic layer was saved and the aqueous phase was extracted three times with ethyl acetate. The extracts were combined with the saved organic layer, washed with 10% sodium bicarbonate, dried over anhydrous magnesium sulfate and concentrated under reduced

pressure to give a first residue. The aqueous phase was acidified and extracted with ethyl acetate. The organic layers were dried and concentrated under reduced pressure to give a second residue. The residues were combined and washed with ethyl acetate to

35

afford the title compound as a white solid, which was identified by NMR and mass spectral analysis. Yield: 0.81 g (42.0%); mp. 213-214°C.


40

EXAMPLES 55-93: Preparation of 3-[5-(N-substituted)sulfamoylcarboxamido-4-chlorophenyl]-1,2,3,4-dihydro-6-(trifluoromethyl)pyrimidin-2,4-diones

Using essentially the above same procedure as described in Example 54 and substituting the appropriate sulfamide starting material, the following compounds were obtained:


45

56



Example No.	x <sup>1</sup>	R <sup>1</sup>	R <sup>2</sup>	R <sup>29</sup>	mp. [°C]
55	F	CH <sub>3</sub>	3-chlorobenzyl	CH <sub>3</sub>	78-79
56	F	H	CH <sub>3</sub>	CH <sub>3</sub>	232-233
57	F	CH <sub>3</sub>	CH <sub>2</sub> -C≡CH	CH <sub>3</sub>	195-196
58	F	H	H	CH <sub>3</sub>	141-142
59	F	CH <sub>3</sub>	allyl	CH <sub>3</sub>	189
60	F	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	81-82
61	F	CH <sub>3</sub>	benzyl	CH <sub>3</sub>	90-91
62	F	C <sub>2</sub> H <sub>5</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	74-76
63	F	H	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	206-207
64	F	H	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	221
65	F	H	C(CH <sub>3</sub> ) <sub>3</sub>	CH <sub>3</sub>	115
66	F	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	215-216
67	F	CH <sub>3</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	72
68	F	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	210-211
69	F	CH <sub>3</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	74
70	F	CH <sub>3</sub>	3-methoxybenzyl	CH <sub>3</sub>	79
71	F	H	CH <sub>2</sub> -C≡CH	CH <sub>3</sub>	195-196
72	F	H	CH <sub>2</sub> -CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	222-223
73	F	CH <sub>3</sub>	phenyl	CH <sub>3</sub>	104-105
74	F	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -phenyl	CH <sub>3</sub>	90-91
75	F	CH <sub>3</sub>	phenyl	NH <sub>2</sub>	120-142
76	F	CH <sub>3</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	NH <sub>2</sub>	117-120
77	F	H	CH <sub>2</sub> -(2-thienyl)	CH <sub>3</sub>	127-128
78	F	H	cyclopentyl	CH <sub>3</sub>	232-233
79	F	H	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	221
80	F	H	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	211
81	F	H	CH <sub>2</sub> -(2-furyl)	CH <sub>3</sub>	178-180
82	F	H	4-methoxybenzyl	CH <sub>3</sub>	186-188
83	F	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	156-157
84	F	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -CN	CH <sub>3</sub>	99-103
85	F	CH <sub>3</sub>	CH <sub>2</sub> -(1,3-dioxolanyl)	CH <sub>3</sub>	93-96

	Example No.	x <sup>1</sup>	R <sup>1</sup>	R <sup>2</sup>	R <sup>29</sup>	mp. [°C]
5	86	F	H	4-chlorobenzyl	CH <sub>3</sub>	95-99
	87	F	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>	CH <sub>3</sub>	126
	88	F		-CH <sub>2</sub> -CH=CH-CH <sub>2</sub> -	CH <sub>3</sub>	231
	89	F	H	cyclopropyl	CH <sub>3</sub>	208
	90	F		-CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -CH <sub>2</sub> -	CH <sub>3</sub>	230
10	91	F	CH <sub>3</sub>	cyclopropyl	CH <sub>3</sub>	156
	92	F	C <sub>2</sub> H <sub>5</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	146
	93	F	H	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	202
	94	F	H	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	227
	95	H	CH <sub>3</sub>	Phenyl	CH <sub>3</sub>	108-110
15	96	F	CH <sub>3</sub>	4-(methoxycarbonyl)phenyl	CH <sub>3</sub>	102
	97	F	C <sub>2</sub> H <sub>5</sub>	Phenyl	CH <sub>3</sub>	214-215
	98	F	CH <sub>3</sub>	3-Pyridyl	CH <sub>3</sub>	208
	99	F	CH <sub>3</sub>	3,4-dichlorophenyl	CH <sub>3</sub>	118
	100	F	CH <sub>3</sub>	3-chlorophenyl	CH <sub>3</sub>	183-184
20	101	F	CH <sub>3</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	93-95
	102	F	CH <sub>3</sub>	CH <sub>3</sub>	NH <sub>2</sub>	252
	103	H	CH <sub>3</sub>	CH <sub>2</sub> -C $\square$ CH	CH <sub>3</sub>	228.2-229.0
	104	F	CH <sub>3</sub>	4-(methoxy)phenyl	CH <sub>3</sub>	136-138
	105	F	CH <sub>3</sub>	4-chlorophenyl	CH <sub>3</sub>	110-111
25	106	F	CH <sub>3</sub>	4-nitrophenyl	CH <sub>3</sub>	111-112
	107	F	CH <sub>3</sub>	4-methylphenyl	CH <sub>3</sub>	102
	108	H	H	H	CH <sub>3</sub>	143.5-145.8
	109	H	CH <sub>3</sub>	CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	187.0-189.5
	110	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	245.5-246.0
30	111	H	CH <sub>3</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	CH <sub>3</sub>	164.1-164.7
	112	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	244.4-245.4
	113	H	CH <sub>3</sub>	CH <sub>2</sub> -CH <sub>2</sub> -C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	167.9-172.0
	114	H	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	228.8-231.5
	115	F	CH <sub>3</sub>	2-methylphenyl	CH <sub>3</sub>	125-127
35	116	F	CH <sub>3</sub>	3-methylphenyl	CH <sub>3</sub>	187-189
	117	F	CH <sub>3</sub>	$\alpha$ -naphthyl	CH <sub>3</sub>	131-133
	118	F	CH <sub>3</sub>	2,4-difluorophenyl	CH <sub>3</sub>	118-119
	119	F	CH <sub>3</sub>	2-chlorophenyl	CH <sub>3</sub>	133
	120	F	CH <sub>3</sub>	2-(trifluoromethyl)phenyl	CH <sub>3</sub>	98-106
40	121	H	CH <sub>3</sub>	4-(phenoxy)phenyl	CH <sub>3</sub>	95
	122	F	CH <sub>3</sub>	4-(trifluoromethyl)phenyl	CH <sub>3</sub>	133
	123	F	CH <sub>3</sub>	4-(dimethylamino)phenyl	CH <sub>3</sub>	87

Example No.	x <sup>1</sup>	R <sup>1</sup>	R <sup>2</sup>	R <sup>29</sup>	mp. [°C]
124	F	CH <sub>3</sub>	4-diphenyl	CH <sub>3</sub>	125-133
125	F	CH <sub>3</sub>	CH(CH <sub>3</sub> )-C <sub>2</sub> H <sub>5</sub>	NH <sub>2</sub>	yellow glass
126	F	CH <sub>3</sub>	C(CH <sub>3</sub> ) <sub>3</sub>	NH <sub>2</sub>	yellow glass
127	F	CH <sub>3</sub>	CH <sub>2</sub> -CH(CH <sub>3</sub> ) <sub>2</sub>	NH <sub>2</sub>	yellow glass
128	F	CH(CH <sub>3</sub> ) <sub>2</sub>	CH(CH <sub>3</sub> ) <sub>2</sub>	NH <sub>2</sub>	yellow glass
129	F	CH <sub>3</sub>	3-(methoxy)phenyl	CH <sub>3</sub>	86
130	H	CH <sub>3</sub>	4-fluorophenyl	CH <sub>3</sub>	120
131	F	CH <sub>3</sub>	3-(dimethylamino)phenyl	CH <sub>3</sub>	85
132	F	CH <sub>3</sub>	3,5-(dichloro)phenyl	CH <sub>3</sub>	104
133	F	CH <sub>3</sub>	CH <sub>2</sub> -CO-OC <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	118-119
134	F	CH <sub>3</sub>			112

Example 135: Preparation of N'-(2-chloro-4-fluoro-5-[4-(di-  
fluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-  
triazol-1-yl]-benzoyl)-N-isopropyl-N-methylsulfamide

Step 1:

2-(2-Fluoro-4-chloro-5-carbomethoxyphenylhydrazonyl)propionic acid

A solution of pyruvic acid (3.92 g, 44.5 mmol) in water (4 ml) was added to a mixture of hydrazine x1 (8.00 g, 36.6 mmol), ethanol (240 ml) and hydrochloric acid (10% strength, 37 ml). The mixture was stirred for 35 minutes at 45-60°C, cooled to 30°C and filtered. The filtrate was concentrated under reduced pressure to ~50% of the original volume and added slowly to water (700 ml). The resultant suspension was stirred for 20 minutes and filtered. The title compound was obtained as a yellow solid. Yield: 9.60 g (90.9%).

Step 2:

2-Chloro-4-fluoro-5-[4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-benzoic acid methyl ester

A mixture of the hydrazone from step 1 (9.00 g, 31.0 mmol) and triethylamine (4.35 ml, 31.0 mmol) was heated to 50°C and treated with a mixture of azide 12 (7.98 g, 29.0 mmol) and toluene (10

ml). The resultant mixture was stirred for 100 minutes at 50°C, cooled to room temperature and concentrated under reduced pressure. The residue was partitioned between water and ethyl acetate. The organic layer was dried and concentrated under  
5 reduced pressure to give the title compound as an off-white solid, which was used in the next step without further purification. Yield: 6.35 g (71.8%).

Step 3:

- 10 2-Chloro-4-fluoro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-benzoic acid methylester

An excess of chlorodifluoromethane (97 g) was bubbled into a mixture of triazolinone from step 2 (7.04 g, 24.6 mmol), tetra-  
15 butylammonium bromide (9.67 g, 30.0 mmol), potassium carbonate (16.6 g, 122 mmol) and dimethylformamide (200 ml) in such a way that  $T < 36^{\circ}\text{C}$  over a period of 30 minutes. The mixture was cooled and filtered. The filtrate was concentrated under reduced  
20 pressure and the residue partitioned between water and methylene chloride. The organic layer was washed with water, dried and concentrated under reduced pressure to give a dark oil (9.00 g), which was chromatographed on silica gel (eluent: ethyl acetate/hexane) to give the title compound. Yield: 1.48 g (17.9%).

25 Step 4:

2-Chloro-4-fluoro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-benzoic acid (IIba)

A mixture of ester II with  $Q = Q^5$ ;  $R^7 = \text{CHF}_2$ ,  $A^4 = \text{O}$ ,  $R^8 = \text{CH}_3$ ,  
30  $X^1 = \text{F}$  and  $X^2 = \text{Cl}$  (0.950 g, 2.83 mmol), acetic acid (10 ml) and hydrochloric acid (6N, 5.0 ml) was stirred for 24 hours at 64-100°C, cooled to room temperature and concentrated under reduced pressure. The residue was partitioned between methylene chloride and water. The organic layer was washed twice with  
35 water, dried and concentrated under reduced pressure to yield the title compound as a pale yellow solid. Yield: 0.42 g (46%); mp.: 132-135°C.

Step 5:

- 40 N'-[[2-chloro-4-fluoro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-benzoyl]]-N-isopropyl-N-methylsulfamide

A mixture of acid II (0.370 g, 1.15 mmol), the sulfamide (0.210  
45 g, 1.38 mmol), carbonyl diimidazole (0.244 g, 1.50 mmol), DBU (0.228 g, 1.50 mmol) and tetrahydrofuran was stirred for four days at ambient temperature. The resultant mixture was treated



with additional sulfamide (0.150 g, 0.986 mmol) and DBU (0.150 g, 0.986 mmol). The mixture was refluxed for two hours, cooled and concentrated under reduced pressure. The residue was taken up in water and acidified with hydrochloric acid (1 N). The mixture was  
5 extracted four times with methylene chloride. The combined extracts were washed with water, dried and concentrated under reduced pressure to give a yellow oil, which was chromatographed on silica gel (eluent: ethyl acetate/hexane), yielding the title compound as an off-white solid. Yield: 0.160 g (30.5%); mp. =  
10 118-122°C.

Example 136: Preparation of N'-(2-chloro-4-fluoro-5-(5,6,7,8-tetrahydro-3-oxo-1,2,4-triazolo[4,3-a]pyridin-2(3H)-yl)-benzoyl)-N,N-dialkylsulfamide

15

Step 1:

2-Chloro-4-fluoro-5-(5,6,7,8-tetrahydro-3-oxo-1,2,4-triazolo[4,3-a]pyridin-2(3H)-yl)-benzoic acid

20 Boron tribromide (19.4 ml, 19.4 mmol, 1N in methylene chloride) was added dropwise to a mixture of I with A = O, X<sup>1</sup> = F, X<sup>2</sup> = Cl, Q = Q<sup>5</sup>, A<sup>4</sup> = O and R<sup>7</sup>+R<sup>8</sup> = -(CH<sub>2</sub>)<sub>4</sub>- (1.90 g, 5.54 mmol) and methylene chloride (20 ml). The resultant mixture was stirred overnight at room temperature. Water (40 ml) was added dropwise  
25 and the mixture stirred for three hours at room temperature. The organic layer was separated and concentrated under reduced pressure to give the title compound as a pale yellow solid. Yield: 1.34 g (77.4%); mp.: 91-94°C.

30


Step 2:  
N'-(2-chloro-4-fluoro-5-(5,6,7,8-tetrahydro-3-oxo-1,2,4-triazolo[4,3-a]pyridin-2(3H)-yl)-benzoyl)-N-isopropyl-N-methylsulfenamide

35 Carbonyl diimidazole (0.500 g, 3.08 mmol) was added to a solution of the compound of step 1 (0.640 g, 2.05 mmol) in anhydrous methylene chloride (15 ml). The resultant mixture was stirred for 30 minutes at room temperature, brought to reflux temperature and immediately cooled to room temperature. The sulfamide (0.370 g,  
40 2.46 mmol) and the DBU (0.470 g, 3.08 mmol) were added and the resultant mixture was stirred overnight at room temperature. The mixture was diluted with ethyl acetate (10 ml) and 3% strength hydrochloric acid (15 ml) and stirred for 10 minutes. Removal of the methylene chloride under reduced pressure gives an aqueous  
45 residue which was extracted with ethyl acetate. The combined organic layers were washed with water and brine, and dried over anhydrous magnesium sulfate. Concentration under reduced pressure

and recrystallization from methylene chloride yielded a solid, which was taken up in methylene chloride and filtered through a column of basic alumina with methanol/methylene chloride, giving the title compound as a white solid. Yield: 0.360 g (39.4%); mp.: 5 250-251°C.

#### Example 137

Using an identical procedure as described in Example 136 hereinabove and the sulfamide  $\text{H}_2\text{N}-\text{SO}_2-\text{N}(\text{CH}_3)-\text{phenyl}$ , the following 10 compound was isolated as a white solid of m.p. 200-201°C:



Example 138: Preparation of N'-[2-chloro-5-(3,5-dimethyl-2,6-dioxo-4-thioxo-1,3,5-triazinan-1-yl)-4-fluorobenzoyl]-N-isopropyl-N-methylsulfamide 20

A solution of 0.502 g (1.45 mmol) of the acid III with  $\text{X}^1 = \text{F}$ ,  $\text{X}^2 = \text{Cl}$ ,  $\text{Q} = \text{Q}^{22}$ ,  $\text{R}^{32} \text{ \& \; } \text{R}^{33} = \text{CH}_3$ ,  $\text{A}^{15} \text{ \& \; } \text{A}^{16} = \text{O}$  and  $\text{A}^{17} = \text{S}$  in 5 ml of tetrahydrofuran was treated with 0.283 g (1.74 mmol) 25 of N,N-carbonyl-diimidazole at 60°C. After this solution had been cooled to ambient temperature, a solution of 0.287 g (1.89 mmol) N,N-methyldiisopropylsulfamide and 0.276 g (1.82 mmol) of DBU in 10 ml of tetrahydrofuran was added and the mixture was stirred overnight. After removal of the volatiles under reduced pressure, 30 the crude product was chromatographed on silica gel with ethyl acetate and cyclohexane. Yield: 0.180 mg of the desired product.

$^1\text{H-NMR}$  (270 MHz; in  $\text{CDCl}_3$ ):  $\delta$  [ppm] = 1.25 (d, 6H), 2.95 (s, 3H,  $\text{SO}_2-\text{NCH}_3$ ), 3.8 (s, 6H,  $\text{NCH}_3$ ), 4.3 (m, 1H), 7.4 (d, 1H, Ar-H), 7.8 35 (d, 1H, Ar-H), 8.9 (bs, 1H, NH).

Example 139: Preparation of N'-[2-chloro-4-fluoro-5-(5-trifluoromethylpyridazon-3-on-2-yl)-benzoyl]-N-isopropyl-N-methylsulfenamides 40

#### Step 1:

2-Fluoro-4-chloro-5-(5-trifluoromethyl-pyridazin-3-on-2-yl)-benzoic acid 45

Pyridazinone ester II with  $\text{X}^1 = \text{F}$ ,  $\text{X}^2 = \text{Cl}$ ,  $\text{Q} = \text{Q}^{27}$ ,  $\text{R}^{34} = \text{H}$ ,  $\text{R}^{35} = \text{CF}_3$  and  $\text{R}^{36} = \text{H}$  (3 g, 8.56 mmol) was dissolved in anhydrous  $\text{CH}_2\text{Cl}_2$

(100 ml). A solution of  $\text{BBr}_3$  (30 ml, 1 M in  $\text{CH}_2\text{Cl}_2$ , 30 mmol) was added to the solution and the mixture was stirred at room temperature for 17 hours. Water (50 ml) was added and the mixture stirred vigorously for 3 hours. A rotary evaporator was used to remove the  $\text{CH}_2\text{Cl}_2$  and the suspended solid was filtered, washed with water and dried to give 2.76 g of the product as a pale yellow solid. Yield: 95%.


Using an identical procedure, the ester II with  $\text{X}^1 = \text{F}$ ,  $\text{X}^2 = \text{Cl}$ ,  $\text{Q} = \text{Q}^{27}$ ,  $\text{R}^{34} = \text{H}$ ,  $\text{R}^{35} = \text{CF}_3$  and  $\text{R}^{36} = \text{CH}_3$ , shown above, was converted to the corresponding acid III in 97% yield.

#### Step 2:

$\text{N}'\text{-}\{2\text{-chloro-4-fluoro-5-(5-trifluoromethyl-pyridazon-3-on-2-yl)}\}$ -benzoyl-N-isopropyl-N-methylsulfenamides

The carboxylic acid III with  $\text{X}^1 = \text{F}$ ,  $\text{X}^2 = \text{Cl}$ ,  $\text{Q} = \text{Q}^{27}$ ,  $\text{R}^{34} = \text{H}$ ,  $\text{R}^{35} = \text{CF}_3$  and  $\text{R}^{36} = \text{CH}_3$  (0.71 g, 2.10 mmol) was dissolved in anhydrous tetrahydrofuran (15 ml) and CDI (0.51 g, 3.15 mmol) was added as a single portion. The mixture was stirred at room temperature for 30 minutes. The mixture was refluxed for 5 minutes, then cooled to room temperature. The sulfamide (0.32 g, 2.10 mmol) was added, followed by DBU (0.48 g, 3.15 mmol). The reaction was stirred at room temperature for 17 hours. 5% strength HCl (15 ml) and ethyl acetate (10 ml) were added to the reaction, and the mixture was stirred vigorously for 10 minutes. The mixture was extracted with ethyl acetate (3 x 15 ml) and the combined extracts were washed with water, dried over  $\text{MgSO}_4$  and concentrated in a rotary evaporator to give a brown semi-solid. The crude product was purified by chromatography on a basic alumina column (eluted with  $\text{CH}_2\text{Cl}_2$ , 1%, 2%  $\text{H}_3\text{C-OH}/\text{CH}_2\text{Cl}_2$  then 1% acetic acid/ $\text{CH}_2\text{Cl}_2$ ) to give the final product Ida.xxx (0.45g) as an off-white solid. Yield: 45%; mp.: 82°C.

The following compounds were prepared using the same procedure and with the appropriate acid and sulfamide.



Example No.	$\text{R}^1$	$\text{R}^2$	$\text{R}^{36}$	mp. [°C]
140	$\text{CH}_3$	phenyl	H	182
141	$\text{CH}_3$	phenyl	$\text{CH}_3$	74-75
142	$\text{CH}_3$	$\text{CH}(\text{CH}_3)_2$	$\text{CH}_3$	181-182

Example No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>36</sup>	mp. [°C]
143	CH <sub>3</sub>	CH <sub>3</sub>	CH <sub>3</sub>	205-206
144	CH <sub>3</sub>	CH <sub>3</sub>	H	184-186

5

Example 145: Preparation of 4-chloro-3-[4-chloro-2-fluoro-5-(N-methyl-N-isopropyl)-sulfamoylcarboxamidophenyl]-5-difluoromethoxy-1-methyl-1H-pyrazole

- 10 A solution of 3-[5-carboxy-4-chloro-2-fluorophenyl]-4-chloro-5-difluoromethoxy-1-methyl-1H-pyrazole (2.00 g, 5.63 mmol) in tetrahydrofuran was treated with N,N'-carbonyldiimidazole (1.13 g, 6.98 mmol), stirred for 1 hour under reflux, cooled to room temperature, treated with N-methyl-N-isopropylsulfamide (1.10 g, 7.23 mmol), stirred for 10 minutes, treated with diazabicyclo-
- 15 undecene (1.06 g, 6.97 mmol), stirred overnight at room temperature and concentrated under reduced pressure. Chromatography on silica gel (cyclohexane/ethylacetate = 4:1) gave 0.90 g of the raw product. Further crystallization from
- 20 cyclohexane/ethylacetate (4:1) yielded 0.45 g (16.4%) of the title compound (analyzed by NMR).  
<sup>1</sup>H-NMR (in CDCl<sub>3</sub>): δ [ppm] = 8.8 (s, 1H, NH), 8.0 (d, 1H), 7.3 (d, 1H), 6.7 (t, 1H), 4.3 (hpt, 1H), 3.8 (s, 3H), 3.0 (s, 3H), 1.2 (d, 6H).

25

Step 1:

2-Chloro-5-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-4-fluorobenzoic acid

- 30 5.3 g (13.4 mmol) of isopropyl ester II with A = O, X<sup>1</sup> = F, X<sup>2</sup> = Cl, X<sup>3</sup> = H, Q = Q<sup>38</sup>, R<sup>40</sup> = Cl, R<sup>41</sup> & R<sup>43</sup> = H and R<sup>42</sup> = CF<sub>3</sub>, dissolved in 25 ml of glacial acid and 125 ml of concentrated HCl were stirred at 70°C for 6 hours and at ambient temperature overnight. Then, the reaction mixture was dripped into ice water
- 35 and the precipitate was filtered off and washed with water. There was obtained 4.1 g as a white solid.  
<sup>1</sup>H-NMR [in (CD<sub>3</sub>)<sub>2</sub>SO]: δ [ppm] = 9.1 (s, 1 H), 8.7 (s, 1 H), 8.1 (d, 1 H), 7.8 (d, 1 H). {Remark: the exchange of the OH proton for those of water resulted in a broad singulett at 3.3 ppm}.

40

Step 2:

N'-[[2-Chloro-5-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-4-fluorobenzoyl]]-N-isopropyl-N-methylsulfamide

- 45 1.0 g (2.8 mmol) free acid from step 1 was dissolved in 10 ml of tetrahydrofuran, 0.57 g (3.5 mmol) of carbonyldiimidazole was added and the mixture was heated to 60°C for 1 hour and cooled to

ambient temperature. Then a mixture of 0.55 g (3.6 mmol) of the sulfamide and 0.54 g (3.5 mmol) of 1,8-diazabicyclo[5,4,0]-undecen-7-ene in 10 ml of tetrahydrofuran was added and stirring was continued at room temperature overnight. The solvent was removed and the crude product was subjected to column chromatography with methyl-tert.-butylether and ethyl acetate. The product-containing fraction was dissolved in methyl-tert.-butylether, washed three times with 10 % strength HCl and twice with water and dried over Na<sub>2</sub>SO<sub>4</sub>. Removal of the solvent gave 0.48 g of the title compound as an oil.

<sup>1</sup>H-NMR [in (CD<sub>3</sub>)<sub>2</sub>SO]: δ [ppm] = 9.1 (s, 1 H), 8.7 (s, 1 H), 7.8-7.7 (m, 2 H), 4.1 (m, 1 H), 2.7 (s, 3 H), 1.1 (m 6 H).

{Remark: the exchange of the N-H protons for those of water resulted in a broad singlett at 3.3 ppm}.

15

Example 146: Preparation of 8-(5'-N-Isopropyl-N-methylsulfamoyl-carboxamido-4'-chloro-2'-fluorophenyl)-4-oxo-7,9-dioxo-1,2,8-triaza(4.3.0.)nonane

5.4 g (45.5 mmol) of thionyl chloride was added to a mixture of 12.0 g (36.4 mmol) acid III with X<sup>1</sup> = F, X<sup>2</sup> = Cl, Q = Q<sup>40</sup>, A<sup>20</sup> & A<sup>21</sup> = O, R<sup>46</sup>+R<sup>47</sup> = -(CH<sub>2</sub>)<sub>3</sub>-O-, and 2 drops of pyridine in 200 ml of 1,2-dichloroethane. After 4 hours at 83°C, the volatiles were removed under reduced pressure and the crude acid chloride (12.6 g) was used without further purification:

0.44 g (2.87 mmol) of N-isopropyl-N-methylaminosulfamide in 50 ml of tetrahydrofuran was added to a suspension of 0.07 g of NaH (97 % purity) in 50 ml of tetrahydrofuran. After 30 minutes at room temperature, 1.0 g (2.87 mmol) of the crude acid chloride was added and the reaction mixture was stirred overnight at ambient temperature and additionally for 2 hours at 50°C. The solvent was removed under reduced pressure, 1 N HCl and methylene chloride were added and the organic layer was separated. Chromatography on silica-gel gave 0.35 g of the title compound; mp.: 115-120°C.

35

#### USE EXAMPLES FOR THE HERBICIDAL ACTIVITY

EXAMPLE 147: Postemergence herbicidal evaluation of test compounds

40

The herbicidal activity of the compounds of the present invention was evaluated by the following tests.

Seedling plants were grown in jiffy flats for about two weeks. The test compounds were dispersed in 80/20 acetone/water mixtures containing 1.0% SUN-IT® II, a methylated seed oil, in sufficient

45

quantities to provide the equivalent of about 0.016 to 0.032 kg per hectare of active compound when applied to the plants through a spray nozzle operating at 40 psi for a predetermined time.

- After spraying, the plants were placed on greenhouse benches and 5 were cared for in accordance with conventional greenhouse procedures. Approximately two to three weeks after treatment, the seedling plants were examined and rated according to the rating scale (0-9) set forth below. Where more than one test is involved for a given compound, the data are averaged.
- 10 Results obtained are reported in Table I below. Where more than one test is involved for a given compound, the data are averaged.

HERBICIDE RATING SCALE		
Rating	% Control as compared to the untreated check	
9	100	
8	91-99	
7	80-90	
6	65-79	
5	45-64	
4	30-44	
3	16-29	
2	6-15	
1	1-5	
0	0	

- The scale is based upon a visual observation of plant stand, vigor, malformation, size, chlorosis and overall plant appearance as compared with a control.

Plant species employed in these evaluations are reported by header abbreviation, common name and scientific name.

PLANT SPECIES EMPLOYED			
Header abbreviation	Common name	Scientific name	
ABUTH	Velvetleaf	Abutilon theophrasti, Medic.	
AMBEL	Ragweed, Common	Ambrosia artemisiifolia, L.	
CHEAL	Lambsquarters	Chenopodium album, L. Common	
IPOHE	Morningglory, Ivy leaf	Ipomoea hederacea, (L) Jacq.	
XANST	Cocklebur	Xanthium strumarium	
ALOMY	Blackgrass	Alopecurus myosuroides	

	DIGSA	Crabgrass, (Hairy) L	Digitaria sanguinalis, (L)Scop
	ECHCG	Barnyardgrass	Echinochloa crus- galli, (L.)Beau
5	SETVI	Green foxtail	Setaria viridis, (L.)Beau
	GLXMA	Soybean	Glycine max, (L.) Merr.
10	TRZAW	Winter wheat	Tritium Aestivum, L. (Winter)
	ZEAMX	Field corn	Zea mays L.

15

20

25

30

35

40

45

Table A  
Postemergence Herbicidal Evaluation

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
53	0.032	9.0	9.0	8.4	9.0	9.0	3.0	6.6	8.8	7.5	8.4	6.2	7.6
	0.016	9.0	8.9	8.1	8.8	9.0	2.3	5.3	8.1	6.4	8.1	5.6	7.2
54	0.032	9.0	9.0	9.0	9.0	9.0	2.0	6.0	8.0	7.0	8.5	6.0	7.5
	0.016	9.0	9.0	9.0	9.0	9.0	2.0	5.0	6.0	7.0	8.0	5.5	6.5
55	0.032	9.0	9.0	7.0	9.0	8.0	2.0	6.0	7.0	8.0	8.0	6.0	8.0
	0.016	9.0	8.0	7.0	8.0	9.0	2.0	5.0	7.0	7.0	8.0	6.0	8.0
56	0.032	9.0	9.0	7.0	9.0	9.0	2.0	5.0	7.0	7.0	8.5	5.0	8.0
	0.016	9.0	8.0	7.0	9.0	9.0	2.0	3.0	7.0	6.0	8.5	4.0	7.5
57	0.032	5.0	2.0	2.0	4.0	3.0	1.0	2.0	2.0	3.0	6.5	2.5	5.0
	0.016	2.0	1.0	1.0	6.0	2.0	0.0	1.0	1.0	1.0	6.0	2.0	2.0
58	0.032	9.0	9.0	9.0	9.0	9.0	3.0	6.0	7.0	8.0	8.5	5.0	8.0
	0.016	9.0	9.0	8.0	9.0	9.0	3.0	5.0	6.0	8.0	8.5	4.5	8.0
59	0.032	9.0	9.0	8.0	9.0	9.0	4.0	5.0	7.0	8.0	8.5	4.5	8.0
	0.016	9.0	7.0	7.0	9.0	9.0	3.0	4.0	6.0	7.0	8.0	4.0	8.0



Table A, continued

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
60	0.032	9.0	9.0	9.0	9.0	9.0	3.0	7.0	7.0	9.0	8.5	6.0	8.0
	0.016	9.0	9.0	8.0	9.0	9.0	3.0	6.0	8.0	8.0	8.5	6.0	8.0
61	0.032	9.0	9.0	8.0	9.0	9.0	1.0	6.0	8.0	9.0	8.5	6.5	7.5
	0.016	9.0	9.0	9.0	8.0	9.0	1.0	7.0	7.0	7.0	8.5	6.0	7.5
62	0.032	9.0	9.0	8.0	9.0	9.0	2.0	5.0	7.0	6.0	8.0	6.5	6.5
	0.016	9.0	9.0	7.0	9.0	9.0	2.0	5.0	6.0	6.0	8.0	5.5	6.5
63	0.032	9.0	9.0	9.0	9.0	9.0	2.0	4.0	6.0	9.0	8.0	7.0	8.0
	0.016	9.0	9.0	7.0	9.0	9.0	2.0	4.0	4.0	7.0	8.0	6.0	7.0
64	0.032	9.0	9.0	9.0	9.0	9.0	5.0	5.0	9.0	9.0	8.0	7.0	8.0
	0.016	9.0	9.0	9.0	9.0	9.0	3.0	5.0	9.0	8.0	7.5	6.0	7.5
65	0.032	9.0	9.0	9.0	9.0	9.0	3.0	8.0	9.0	8.0	8.5	6.5	8.0
	0.016	9.0	9.0	8.0	9.0	9.0	2.0	6.0	8.0	8.0	8.0	5.5	7.5
66	0.032	9.0	9.0	9.0	9.0	9.0	2.0	6.0	8.0	9.0	9.0	6.5	7.5
	0.016	9.0	9.0	8.0	9.0	9.0	1.0	5.0	8.0	8.0	8.0	5.0	7.0
67	0.032	9.0	9.0	9.0	9.0	9.0	3.0	6.0	8.0	8.0	8.5	7.0	7.5
	0.016	9.0	9.0	9.0	9.0	9.0	2.0	6.0	8.0	8.0	8.0	6.5	7.5

Table A, continued

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
68	0.032	9.0	9.0	8.0	9.0	9.0	1.0	6.0	8.0	9.0	8.5	5.5	7.5
	0.016	9.0	9.0	8.0	9.0	9.0	1.0	6.0	8.0	7.0	8.5	5.0	7.0
69	0.032	9.0	9.0	8.0	9.0	9.0	2.0	6.0	7.0	6.0	8.0	6.5	7.5
	0.016	9.0	9.0	8.0	9.0	9.0	1.0	4.0	5.0	5.0	8.0	5.0	7.0
70	0.032	9.0	7.0	7.0	9.0	-	1.0	3.0	8.0	5.0	7.5	5.0	7.0
	0.016	9.0	5.0	6.0	9.0	-	1.0	2.0	4.0	4.0	6.5	4.5	7.0
71	0.032	9.0	9.0	5.0	9.0	-	1.0	3.0	3.0	5.0	8.0	4.5	6.5
	0.016	9.0	6.0	5.0	8.0	-	1.0	2.0	2.0	4.0	8.0	4.5	6.5
72	0.032	9.0	9.0	8.0	9.0	-	2.0	4.0	8.0	6.0	8.5	6.5	7.0
	0.016	9.0	9.0	8.0	9.0	-	1.0	4.0	6.0	5.0	8.5	5.5	7.0
73	0.032	9.0	9.0	9.0	9.0	-	3.0	6.0	7.0	5.0	8.5	6.5	7.5
	0.016	9.0	9.0	8.0	9.0	-	2.0	4.0	4.0	4.0	8.5	5.5	7.0
76	0.032	9.0	8.0	6.0	9.0	9.0	1.0	4.0	6.0	6.0	7.5	6.0	6.5
	0.016	9.0	8.0	6.0	9.0	9.0	1.0	3.0	5.0	5.0	7.5	5.0	5.5
77	0.032	9.0	8.0	7.0	9.0	9.0	1.0	4.0	6.0	6.0	7.5	5.0	6.5
	0.016	9.0	8.0	6.0	9.0	9.0	1.0	4.0	4.0	5.0	7.5	4.5	6.0

Table A, continued

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
78	0.032	9.0	9.0	9.0	9.0	9.0	2.0	6.0	9.0	9.0	8.0	6.5	7.0
	0.016	9.0	9.0	7.0	9.0	9.0	2.0	5.0	8.0	8.0	7.0	5.5	6.5
80	0.032	9.0	8.0	4.0	9.0	9.0	1.0	4.0	6.0	6.0	8.0	4.0	4.0
	0.016	7.0	7.0	3.0	9.0	8.0	0.0	3.0	4.0	5.0	8.0	3.0	1.5
82	0.032	9.0	9.0	9.0	9.0	9.0	2.0	3.0	8.0	7.0	8.5	6.0	6.5
	0.016	9.0	9.0	9.0	9.0	9.0	1.0	3.0	7.0	6.0	8.0	5.5	6.5
83	0.032	9.0	8.0	5.0	9.0	9.0	1.0	4.0	8.0	8.0	7.5	5.5	6.0
	0.016	9.0	6.0	5.0	8.0	9.0	0.0	3.0	8.0	7.0	7.0	4.5	5.5
84	0.032	9.0	9.0	9.0	9.0	9.0	2.0	6.0	9.0	8.0	8.5	5.5	8.0
	0.016	9.0	8.0	7.0	9.0	9.0	1.0	5.0	8.0	7.0	7.0	5.0	7.0

**EXAMPLE 148: Preemergence herbicidal evaluation Of test compounds**

The herbicidal activity of the compounds of the present invention was evaluated by the following tests wherein the seeds of a variety of monocotyledonous and dicotyledonous plants were separately mixed with potting soil and planted on top of approximately one inch of soil in separate pint cups. After planting, the cups were sprayed with the selected aqueous acetone solution containing test compound in sufficient quantity to provide the equivalent of about 0.125 to 0.250 kg per hectare of test compound per cup. The treated cups were then placed on greenhouse benches, watered and cared for in accordance with conventional greenhouse procedures. From two to four weeks after treatment, the tests were terminated and each cup was examined and rated according to the rating system provided in Example 94. When more than one test was performed for a given compound, the data were averaged. The results obtained are shown in Table B.

Table B  
Preemergence Herbicidal Evaluation

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
53	0.25	9.0	9.0	9.0	9.0	9.0	2.7	8.8	8.4	7.3	8.8	2.0	4.3
	0.125	9.0	8.9	9.0	9.0	9.0	1.3	8.1	7.5	5.7	8.1	1.3	2.3
54	0.25	9.0	8.0	9.0	1.0	3.0	0.0	6.0	3.0	3.0	0.0	0.0	0.0
	0.125	9.0	7.0	9.0	0.0	1.0	0.0	4.0	3.0	2.0	0.0	0.0	0.0
55	0.25	9.0	9.0	9.0	9.0	7.0	0.0	8.0	9.0	8.0	8.0	2.0	1.0
	0.125	9.0	8.0	8.0	8.0	6.0	0.0	8.0	5.0	6.0	8.0	1.0	0.0
56	0.25	9.0	9.0	9.0	9.0	9.0	0.0	7.0	7.0	8.0	9.0	0.0	2.0
	0.125	9.0	9.0	9.0	8.0	8.0	0.0	7.0	3.0	4.0	2.0	0.0	0.0
57	0.25	2.0	2.0	8.0	3.0	0.0	0.0	3.0	2.0	1.0	1.0	0.0	0.0
	0.125	2.0	2.0	6.0	0.0	0.0	0.0	1.0	1.0	0.0	0.0	0.0	0.0
58	0.25	9.0	9.0	9.0	9.0	8.0	0.0	3.0	6.0	6.0	9.0	3.0	1.0
	0.125	9.0	9.0	9.0	9.0	5.0	0.0	0.0	4.0	2.0	9.0	0.0	0.0
59	0.25	9.0	9.0	9.0	9.0	9.0	4.0	7.0	3.0	9.0	8.0	0.0	2.0
	0.125	9.0	9.0	9.0	9.0	0.0	0.0	3.0	0.0	4.0	2.0	0.0	0.0

Table B, continued

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	KANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
60	0.25	9.0	9.0	9.0	6.0	4.0	0.0	6.0	4.0	9.0	6.0	0.0	0.0
	0.125	9.0	9.0	9.0	7.0	0.0	0.0	6.0	0.0	5.0	0.0	0.0	0.0
61	0.25	9.0	9.0	9.0	9.0	9.0	0.0	9.0	7.0	6.0	9.0	3.0	2.0
	0.125	9.0	9.0	9.0	9.0	9.0	0.0	6.0	4.0	3.0	9.0	2.0	2.0
62	0.25	9.0	9.0	9.0	9.0	9.0	0.0	7.0	6.0	7.0	9.0	5.0	1.0
	0.125	9.0	9.0	9.0	9.0	9.0	0.0	5.0	3.0	3.0	5.0	3.0	1.0
63	0.25	9.0	9.0	9.0	9.0	-	0.0	8.0	9.0	8.0	5.0	0.0	0.0
	0.125	9.0	9.0	9.0	9.0	-	0.0	3.0	4.0	4.0	1.0	0.0	0.0
64	0.25	9.0	9.0	9.0	9.0	-	4.0	9.0	7.0	9.0	-	3.0	2.0
	0.125	9.0	9.0	9.0	9.0	-	0.0	8.0	3.0	8.0	-	0.0	0.0
65	0.25	9.0	9.0	9.0	9.0	-	0.0	9.0	9.0	6.0	-	1.0	1.0
	0.125	9.0	9.0	9.0	9.0	-	0.0	7.0	8.0	3.0	-	1.0	0.0
66	0.25	9.0	9.0	9.0	9.0	-	2.0	8.0	8.0	6.0	-	1.0	2.0
	0.125	9.0	9.0	9.0	9.0	-	0.0	4.0	4.0	4.0	0.0	1.0	1.0
67	0.25	9.0	7.0	9.0	9.0	-	0.0	8.0	9.0	8.0	-	1.0	1.0
	0.125	9.0	9.0	9.0	9.0	-	0.0	7.0	8.0	8.0	-	0.0	0.0

Table B, continued

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
68	0.25	9.0	9.0	8.0	9.0	9.0	0.0	4.0	4.0	8.0	8.0	0.0	2.0
	0.125	9.0	9.0	8.0	9.0	9.0	0.0	3.0	3.0	4.0	7.0	0.0	1.0
69	0.25	9.0	9.0	9.0	9.0	4.0	0.0	3.0	3.0	1.0	3.0	0.0	3.0
	0.125	9.0	8.0	9.0	9.0	3.0	0.0	3.0	2.0	0.0	0.0	0.0	2.0
70	0.25	9.0	9.0	9.0	9.0	9.0	0.0	5.0	7.0	5.0	3.0	1.0	0.0
	0.125	9.0	9.0	8.0	9.0	8.0	0.0	2.0	2.0	3.0	1.0	0.0	0.0
71	0.25	9.0	8.0	7.0	9.0	7.0	0.0	3.0	3.0	2.0	2.0	0.0	1.0
	0.125	9.0	8.0	9.0	9.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
72	0.25	9.0	9.0	9.0	9.0	9.0	2.0	7.0	6.0	8.0	9.0	1.0	0.0
	0.125	9.0	9.0	9.0	9.0	7.0	1.0	2.0	7.0	4.0	8.0	0.0	0.0
73	0.25	9.0	9.0	9.0	9.0	9.0	0.0	3.0	3.0	5.0	0.0	0.0	0.0
	0.125	9.0	9.0	9.0	9.0	3.0	0.0	3.0	0.0	2.0	0.0	0.0	0.0
76	0.25	9.0	9.0	9.0	9.0	5.0	2.0	3.0	2.0	2.0	3.0	1.0	1.0
	0.125	9.0	8.0	8.0	5.0	3.0	0.0	2.0	1.0	1.0	1.0	0.0	1.0
77	0.25	9.0	9.0	9.0	9.0	8.0	0.0	4.0	3.0	2.0	-	0.0	1.0
	0.125	9.0	9.0	9.0	7.0	3.0	0.0	2.0	2.0	2.0	1.0	0.0	0.0

Table B, continued

Ex. No.	Rate [kg/ha]	ABUTH	AMBEL	CHEAL	IPOHE	XANST	ALMOY	DIGSA	ECHCG	SETVI	GLXMA	TRZAW	ZEAMX
78	0.25	9.0	9.0	9.0	9.0	9.0	2.0	8.0	7.0	7.0	5.0	1.0	1.0
	0.125	9.0	9.0	9.0	9.0	9.0	1.0	4.0	4.0	5.0	3.0	1.0	0.0
80	0.25	5.0	3.0	8.0	0.0	1.0	0.0	1.0	3.0	0.0	0.0	0.0	0.0
	0.125	1.0	1.0	5.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
82	0.25	9.0	9.0	9.0	9.0	9.0	0.0	4.0	4.0	7.0	9.0	1.0	2.0
	0.125	9.0	9.0	9.0	9.0	9.0	0.0	4.0	3.0	3.0	9.0	0.0	0.0
83	0.25	9.0	8.0	9.0	8.0	5.0	0.0	9.0	5.0	6.0	4.0	0.0	2.0
	0.125	8.0	8.0	9.0	7.0	4.0	0.0	1.0	4.0	3.0	1.0	0.0	0.0
84	0.25	9.0	9.0	9.0	9.0	9.0	1.0	8.0	6.0	7.0	7.0	2.0	1.0
	0.125	9.0	9.0	9.0	9.0	9.0	0.0	7.0	5.0	7.0	2.0	0.0	0.0



## EXAMPLE 149

The herbicidal action of the uracil substituted phenyl sulfamoyl carboxamides no. Ij.86, Ip.86 and Iy.86 was demonstrated by the following greenhouse experiments:

The culture containers used were plastic flowerpots containing loamy sand with approximately 3.0% of humus as substrate. The seeds of the test plants were sown separately for each species.

For the post-emergence treatment, the test plants were grown to a plant height of from 3 to 15 cm, depending on the plant habit, and only then treated with the active ingredients which had been suspended or emulsified in water. To this end, the test plants were either sown directly and grown in the same containers, or they were first grown separately as seedlings and transplanted into the test containers a few days prior to treatment. The rate of application for the post-emergence treatment was 15.6 or 7.8 g/ha active ingredient.

Depending on the species, the plants were kept at from 10 - 25°C and 20 - 35°C, respectively. The test period extended over 2 to 4 weeks. During this time, the plants were tended, and their response to the individual treatments was evaluated.

Evaluation was carried out using a scale of from 0 to 100. 100 means no emergence of the plants, or complete destruction of at least the aerial parts, and 0 means no damage or normal course of growth.

The plants used in the greenhouse experiments belonged to the following species:

Scientific Name	Common Name
<i>Amaranthus retroflexus</i> (AMARE)	Redroot pigweed
<i>Pharbitis purpurea</i> (PHBPU)	common or tall morningglory
<i>Polygonum persicaria</i> (POLPE)	redshank; ladysthumb

At a rate of application of 15.6 or 7.8 g/ha of a.i., compound nos. Ij.86, Ip.86 and Iy.86 showed a very good herbicidal action against the abovementioned undesired plants.

## DESICCANT/DEFOLIANT ACTIVITY OF THE COMPOUNDS I

## EXAMPLE 150: greenhouse-trials

The test plants used were young cotton plants with 4 leaves (without cotyledons) which had been grown under greenhouse conditions (relative atmospheric humidity 50 to 70%; day/night temperature 27/20°C).

The young cotton plants were subjected to foliar treatment to run-off point with aqueous preparations of the active ingredients (with an addition of 0.15% by weight of the fatty alcohol alkoxylate Plurafac® LF 700<sup>1)</sup>, based on the spray mixture). The amount of water applied was 1000 l/ha (converted). After 13 days, the number of leaves shed and the degree of defoliation in % were determined.

No leaves were shed in the untreated control plants.

1) a low-foam, nonionic surfactant from BASF AG

#### EXAMPLE 151: field trials

Field evaluations of preharvest desiccant activity were performed at several different locations using compound Ia.86.

In each experiment treatments was replicated three times in a randomized complete block experimental design. Potatoes were grown using good agronomic practices of each area. Treatments were applied a few weeks before planned potato harvest.


The test compound was formulated as an emulsifiable concentrate (EC) formulation with 120 grams a.i./liter. The formulation was diluted with water, spray adjuvants were added, and the treatment solution was applied to the foliage of potatoes in from 187 to 600 l/ha of total spray volume. Unless indicated otherwise, the treatments also contained 15 v/v of methylated seed oil adjuvant (Hasten or SUN-IT II. In the case of split applications, the second application was made about 1 week after the initial application.

At various intervals after treatment, desiccation of stems and leaves was evaluated separately, on a visual % desiccation scale. In each test the test compound was compared to appropriate commercial standards at the normal rate for each standard for the area.

The results showed that the abovementioned compound is very effective to desiccate the leaves and stems of potato plants.

## Claims:

1. Uracil substituted phenyl sulfamoyl carboxamides of the formula I



wherein the variables have the following meanings:

A oxygen or sulfur;

X<sup>1</sup> hydrogen, halogen or C<sub>1</sub>-C<sub>4</sub>-alkyl;

X<sup>2</sup> hydrogen, cyano, CS-NH<sub>2</sub>, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl or C<sub>1</sub>-C<sub>4</sub>-haloalkyl;

X<sup>3</sup> hydrogen, cyano, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxy-alkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>3</sub>-C<sub>6</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl or optionally substituted benzyl;


R<sup>1</sup> and R<sup>2</sup> independently of one another

hydrogen, halogen, OR<sup>48</sup>, C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, C<sub>3</sub>-C<sub>10</sub>-alkynyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, phenyl, benzyl or C<sub>5</sub>-C<sub>7</sub>-cycloalkenyl, whereas each of the last-mentioned 7 groups can be substituted with any combination of one to six halogen atoms, one to three C<sub>1</sub>-C<sub>6</sub>-alkoxy groups, one or two C<sub>1</sub>-C<sub>8</sub>-haloalkoxy groups, one or two cyano groups, one or two C<sub>3</sub>-C<sub>7</sub>-cycloalkyl groups, one or two C(O)R<sup>49</sup> groups, one or two CO-OR<sup>50</sup> groups, one or two CO-SR<sup>51</sup> groups, one or two CO-NR<sup>52</sup>R<sup>53</sup> groups, one to three OR<sup>54</sup> groups, one to three SR<sup>54</sup> groups, one optionally substituted four to 10-membered monocyclic or fused bicyclic heterocyclic ring, one or two optionally substituted phenyl groups or one or two optionally substituted benzyl groups,


or R<sup>1</sup> and R<sup>2</sup> together with the atom to which they are attached form a 3- to 7-membered heterocyclic ring;

Q is selected from


5




Q2




Q3




Q4




10




Q6




Q7




Q8




15




Q10




Q11




Q12




25




Q14



Q15




Q16




30


35




Q18



Q19




Q20



40

45




wherein

A<sup>1</sup> to A<sup>17</sup> are each independently oxygen or sulfur;

- 5  $R^3$ ,  $R^4$ ,  $R^7$ ,  $R^8$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{18}$ ,  $R^{19}$ ,  $R^{27}$ ,  $R^{29}$ ,  $R^{32}$ ,  $R^{33}$ ,  $R^{38}$ ,  $R^{39}$ ,  $R^{44}$ ,  $R^{45}$ ,  $R^{46}$  and  $R^{47}$  are each independently hydrogen, cyano, amino,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -haloalkoxy,  $C_3$ - $C_7$ -cycloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_3$ - $C_6$ -alkynyl, benzyl,  $OR^{55}$ ,  $C_1$ - $C_3$ -cyanoalkyl, or
- 10  $R^3$  and  $R^4$ ,  $R^7$  and  $R^8$ ,  $R^{11}$  and  $R^{12}$ ,  $R^{18}$  and  $R^{19}$  or  $R^{46}$  and  $R^{47}$  may be taken together with the atoms to which they are attached to represent a four- to seven-membered ring, optionally interrupted by oxygen, sulfur or nitrogen and optionally substituted with one or more halogen or  $C_1$ - $C_4$ -alkyl groups;
- 15  $R^5$ ,  $R^6$ ,  $R^9$ ,  $R^{10}$ ,  $R^{15}$ ,  $R^{16}$ ,  $R^{20}$ ,  $R^{21}$ ,  $R^{30}$ ,  $R^{31}$ ,  $R^{35}$ ,  $R^{36}$ ,  $R^{41}$ ,  $R^{42}$  and  $R^{43}$  are each independently hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_3$ - $C_7$ -cycloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_3$ - $C_6$ -alkynyl,  $OR^{56}$ ,  $S(O)_nR^{57}$ ,  $O-SO_2-R^{57}$ ,  $NR^{58}R^{59}$  or
- 20  $R^5$  and  $R^6$ ,  $R^9$  and  $R^{10}$ ,  $R^{15}$  and  $R^{16}$ ,  $R^{20}$  and  $R^{21}$  or  $R^{30}$  and  $R^{31}$  may be taken together with the atoms to which they are attached to represent a four- to seven membered ring optionally substituted with one or more halogen or  $C_1$ - $C_4$ -alkyl groups;
- 25  $R^{13}$ ,  $R^{14}$ ,  $R^{22}$ ,  $R^{23}$ ,  $R^{25}$  and  $R^{26}$  are each independently hydrogen, halogen or  $C_1$ - $C_6$ -alkyl;
- 30  $R^{17}$ ,  $R^{28}$ ,  $R^{34}$ ,  $R^{37}$  or  $R^{40}$  are each independently hydrogen, halogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_3$ - $C_7$ -cycloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -haloalkenyl,  $C_3$ - $C_6$ -alkynyl,  $OR^{60}$  or  $SR^{61}$ ;
- 35  $R^{24}$  is hydrogen,  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -haloalkyl,  $C_2$ - $C_4$ -alkenyl,  $C_3$ - $C_4$ -alkynyl,  $C_1$ - $C_4$ -haloalkoxy or amino;
- 40  $R^{48}$ ,  $R^{49}$ ,  $R^{50}$ ,  $R^{51}$ ,  $R^{52}$ ,  $R^{53}$ ,  $R^{54}$ ,  $R^{55}$ ,  $R^{56}$ ,  $R^{57}$ ,  $R^{58}$ ,  $R^{59}$ ,  $R^{60}$  and  $R^{61}$  are independently of one another hydrogen,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_3$ - $C_7$ -cycloalkyl,  $C_2$ - $C_6$ -alkenyl,  $C_3$ - $C_6$ -alkynyl, optionally substituted phenyl or optionally substituted benzyl;
- 45  $n$  is zero, 1 or 2;
- and the agriculturally useful salts of the compounds I.

2. The compound defined in claim 1, wherein Q is selected from Q<sup>5</sup>, Q<sup>7</sup>, Q<sup>21</sup>, Q<sup>22</sup>, Q<sup>27</sup>, Q<sup>32</sup>, Q<sup>38</sup>, Q<sup>39</sup> and Q<sup>40</sup>.
3. 2-Chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-(dimethylsulfamoyl)-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(ethyl)propylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-(sec.-butylsulfamoyl)-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-(tert.-butylsulfamoyl)-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(ethyl)methylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(methyl)propylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-(diethyl)sulfamoyl)-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-amino-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(methyl)isopropylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(sec.-butyl)methylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(methyl)phenylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(methyl)isopropylsulfamoyl]-4-fluorobenzamide,  
2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2H)-pyrimidinyl]-N-[(n-butyl)methylsulfamoyl]-4-fluorobenzamide.
- N'-[[2-chloro-4-fluoro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-benzoyl]]-N-isopropyl-N-methylsulfamide,  
N'-{2-chloro-4-fluoro-5-(5,6,7,8-tetrahydro-3-oxo-1,2,4-triazolo[4,3-a]pyridin-2(3H)-yl)-benzoyl}-N-isopropyl-N-methylsulfenamide,  
N'-[2-chloro-5-(3,5-dimethyl-2,6-dioxo-4-thioxo-1,3,5-triazinan-1-yl)-4-fluorobenzoyl]-N-isopropyl-N-methylsulfamide,

- N'-[2-chloro-4-fluoro-5-(5-trifluoromethylpyridazon-3-on-2-yl)-benzoyl]-N-isopropyl-N-methylsulfenamide,  
4-chloro-3-[4-chloro-2-fluoro-5-(N-methyl-N-isopropyl)-sulfamoylcarboxamidophenyl]-5-difluoromethoxy-1-methyl-  
1H-pyrazole,  
N'-[[2-chloro-5-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-4-fluorobenzoyl]]-N-isopropyl-N-methylsulfamide and  
8-(5'-N-isopropyl-N-methylsulfamoylcarboxamido-4'-chloro-2'-fluorophenyl)-4-oxo-7,9-dioxo-1,2,8-triaza(4.3.0.)nonane.
4. A herbicidal composition, comprising a herbicidally effective amount of at least one compound of the formula I or of an agriculturally useful salt of I as defined in claim 1, and at least one inert liquid and/or solid carrier and, if desired, at least one surfactant.
5. A composition for the desiccation and/or defoliation of plants, comprising such an amount of at least one compound of the formula I or of an agriculturally useful salt of I, as defined in claim I, that it acts as a desiccant and/or defoliant, and at least one inert liquid and/or solid carrier and, if desired, at least one surfactant.
6. A method of controlling undesirable vegetation, which comprises allowing a herbicidally active amount of at least one compound of the formula I or of an agriculturally useful salt of I, as defined in claim 1, to act on plants, their environment or on seed.
7. A method for desiccation/defoliation of plants, which comprises allowing such an amount of at least one compound of the formula I or an agriculturally useful salt of I, as defined in claim 1, to act on plants that it acts as a defoliant and/or desiccant.
8. A method as claimed in claim 8, wherein cotton is treated.
9. A process for the preparation of a compound of formula I as defined in claim 1, which process comprises reacting a benzoic acid derivative of the formula II




II,

wherein Q, X<sup>1</sup> and X<sup>2</sup> are as defined in claim 1,




optionally in the presence of a coupling agent,  
or the corresponding acid chloride of II,  
with a sulfamide of the formula III




wherein  $\text{X}^3$ ,  $\text{R}^1$  and  $\text{R}^2$  are as defined in claim 1.

- 10 10. A process for the preparation of a compound of formula I as defined in claim 1, where A is oxygen,  $\text{X}^3$  is hydrogen, Q is  $\text{Q}^{21}$ ,  $\text{A}^8$  &  $\text{A}^9$  are oxygen and  $\text{R}^{29}$  is hydrogen, which process comprises reacting an aniline intermediate VI




- 20 wherein  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{R}^1$  and  $\text{R}^2$  are as defined in claim 1, with an oxazinone compound of the formula VII




optionally followed by alkylation and hydrolysis.


- 30 11. A process for the preparation of a compound of formula VI



- wherein  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{X}^3$ ,  $\text{R}^1$  and  $\text{R}^2$  are as defined in claim 1, which process comprises treating a sulfamoyl carboxamide X



- 45 with a nitration reagent to give the corresponding nitrated compound XI




XI,

5

and subsequently reducing the nitro group with a transition metal under acidic conditions or with a complex hydride.


10 12. A process for the preparation of a sulfamoyl carboxamide X



X,

15


wherein  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{X}^3$ ,  $\text{R}^1$  and  $\text{R}^2$  are as defined in claim 1, which process comprises reacting a benzoic acid IX



IX,

20

25 optionally in the presence of a coupling agent, or the corresponding acid chloride of IX, with a sulfamide of the formula III



III.

30

35

40

45

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number  
**WO 01/83459 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 239/54**,  
249/12, 471/04, 237/14, 231/20, 213/61, A01N 43/54

(21) International Application Number: **PCT/EP01/04850**

(22) International Filing Date: **30 April 2001 (30.04.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
60/201,824 4 May 2000 (04.05.2000) **US**

(71) Applicant: **BASF AKTIENGESELLSCHAFT**  
[DE/DE]: 67056 Ludwigshafen (DE).

(72) Inventors: **CARLSEN, Marianne**; 222 Arborlea Avenue, Yardley, PA 19067 (US). **GUACIARO, Michael, Anthony**; 21 Inverness Lane, East Windsor, NJ 08520 (US). **TAKASUGI, James, Jan**; 11 Pembroke Court, Lawrenceville, NJ 08648 (US).

(74) Common Representative: **BASF AKTIENGESELLSCHAFT**; 67056 Ludwigshafen (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


**Published:**

— with international search report

(88) Date of publication of the international search report:  
16 May 2002

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: URACIL SUBSTITUTED PHENYL SULFAMOYL CARBOXAMIDES



(57) Abstract: Novel uracil substituted phenyl sulfamoyl carboxamides (I) and salts thereof, where A = oxygen or sulfur; X<sup>1</sup> = H, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl; X<sup>2</sup> = H, CN, CS-NH<sub>2</sub>, halogen, C<sub>1</sub>-C<sub>4</sub>-alkyl, C<sub>1</sub>-C<sub>4</sub>-haloalkyl; X<sup>3</sup> = H, CN, C<sub>1</sub>-C<sub>6</sub>-alkyl, C<sub>1</sub>-C<sub>6</sub>-alkoxyalkyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, C<sub>3</sub>-C<sub>8</sub>-alkenyl, C<sub>3</sub>-C<sub>6</sub>-alkynyl, optionally substituted benzyl; R<sup>1</sup>, R<sup>2</sup> = H, halogen, optionally substituted hydroxy, C<sub>1</sub>-C<sub>10</sub>-alkyl, C<sub>2</sub>-C<sub>10</sub>-alkenyl, C<sub>3</sub>-C<sub>10</sub>-alkynyl, C<sub>3</sub>-C<sub>7</sub>-cycloalkyl, phenyl, benzyl or C<sub>5</sub>-C<sub>7</sub>-cycloalkenyl, or R<sup>1</sup> + R<sup>2</sup> together with the atom to which they are attached form a 3- to 7-membered heterocyclic ring; Q is selected from Q<sup>1</sup> to Q<sup>40</sup> as defined in the description. Use: as herbicides; for the desiccation/defoliation of plants.

WO 01/83459 A3

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 01/04850

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/54 C07D249/12 C07D471/04 C07D237/14 C07D231/20  
C07D213/61 A01N43/54

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DE 197 31 784 A (BAYER) 4 February 1999 (1999-02-04) claims	1,4-9
A	DE 196 10 786 A (BAYER) 18 September 1997 (1997-09-18) claims	1,4-9
A	DE 196 27 901 A (BAYER) 15 January 1998 (1998-01-15) claims	1,4-9
A	DE 196 32 005 A (BAYER) 12 February 1998 (1998-02-12) cited in the application claims	1,4-9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

25 January 2002

Date of mailing of the international search report

06/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Francois, J

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/04850

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 19731784	A	04-02-1999	DE 19731784 A1	04-02-1999
			AU 8860898 A	16-02-1999
			WO 9905125 A1	04-02-1999
DE 19610786	A	18-09-1997	DE 19610786 A1	18-09-1997
			AU 2094997 A	01-10-1997
			BR 9708186 A	27-07-1999
			CZ 9802930 A3	13-01-1999
			WO 9733876 A1	18-09-1997
			EP 0888314 A1	07-01-1999
			HU 9902109 A2	29-11-1999
			JP 2000506177 T	23-05-2000
DE 19627901	A	15-01-1998	DE 19627901 A1	15-01-1998
			AU 3441297 A	09-02-1998
			BR 9710238 A	10-08-1999
			CA 2260201 A1	22-01-1998
			WO 9802422 A1	22-01-1998
			EP 0912527 A1	06-05-1999
			JP 2000514805 T	07-11-2000
DE 19632005	A	12-02-1998	DE 19632005 A1	12-02-1998
			AU 718868 B2	20-04-2000
			AU 3771497 A	06-03-1998
			BR 9711033 A	17-08-1999
			CN 1227548 A	01-09-1999
			WO 9806706 A1	19-02-1998
			EP 0923562 A1	23-06-1999
			JP 2001503021 T	06-03-2001
			US 6130225 A	10-10-2000

**THIS PAGE BLANK (USPTO)**